

**LEMBAR  
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW  
KARYA ILMIAH : PROSIDING**

Judul Karya Ilmiah : **Production and Quality of Seven Basil (*Ocimum basilicum* L.) Accessions in Various Composition of Urea Fertilizer and Mexican Sunflower Compost**

Jumlah Penulis : 4 Orang

Status Pengusul : Penulis Pertama

Identitas Prosiding : a. Nama Prosiding : Proceedings of The International Seminar on Promoting Local Resources for Sustainable Agriculture and Development  
b. ISBN/ISSN : 978-94-6239-391-2/2468-7547  
c. Tahun Terbit, Lokasi : 11 Juni 2021  
d. Halaman/Penerbit : 267-273 / Atlantis Press  
e. Repository/Web : <https://www.atlantis-press.com/proceedings/isplrsad-20/125957666>  
f. Terindeks di : CPCI, SCIE, ESCI, Google Scholar, Scopus

Kategori Publikasi Karya Ilmiah :  *Prosiding* Forum Ilmiah Internasional  
(beri ✓ pada kategori yang tepat)  *Prosiding* Forum Ilmiah Nasional

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal <i>Prosiding</i>		Nilai Akhir Yang Diperoleh
	Internasional <b>25</b>	Nasional <input type="text"/>	
Kelengkapan Unsur Isi Buku (10%)	2.5		2.5
Ruang Lingkup & Kedalaman Pembahasan (30%)	7.5		7.5
Kecukupan & Kemutakhiran Data/Informasi & Metodologi (30%)	7.5		7.5
Kelengkapan Unsur & Kualitas Terbitan/Prosiding (30%)	7.5		7.5
Total = (100%)	25		25

**CATATAN PENILAIAN ARTIKEL OLEH REVIEWER:**

1. Tentang Kelengkapan dan Kesesuaian Unsur Isi Prosiding :

*Isi prosiding lengkap dan sesuai dengan artikel*

2. Tentang Ruang Lingkup dan Kedalaman Pembahasan :

*Ruang lingkup dan pembahasan sangat baik*

3. Kecukupan & Kemutakhiran Data/Informasi & Metodologi :

*Sumber referensi cukup mutakhir*

4. Kelengkapan Unsur & Kualitas Terbitan :

*Penyajiannya melalui seminar internasional*

Reviewer 1,



**Dr. Ir. Anggraeni, M.Si**

NIK/NIDN : 0006076301

Jabatan Fungsional/Pangkat : Lektor Kepala, Pembina IV/a  
Unit Kerja : Universitas Djuanda

**LEMBAR  
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW  
KARYA ILMIAH : PROSIDING**

Judul Karya Ilmiah : **Production and Quality of Seven Basil (*Ocimum basilicum* L.) Accessions in Various Composition of Urea Fertilizer and Mexican Sunflower Compost**

Jumlah Penulis : 4 Orang

Status Pengusul : Penulis Pertama

Identitas Prosiding : a. Nama Prosiding : Proceedings of The International Seminar on Promoting Local Resources for Sustainable Agriculture and Development  
b. ISBN/ISSN : 978-94-6239-391-2/2468-7547  
c. Tahun Terbit, Lokasi : 11 Juni 2021  
d. Halaman/Penerbit : 267-273 / Atlantis Press  
e. Repository/Web : <https://www.atlantispress.com/proceedings/isplrsad-20/125957666>  
f. Terindeks di : CPCI, SCIE, ESCI, Google Scholar, Scopus

Kategori Publikasi Karya Ilmiah :  Prosiding Forum Ilmiah Internasional  
(beri ✓ pada kategori yang tepat)  Prosiding Forum Ilmiah Nasional

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Prosiding		Nilai Akhir Yang Diperoleh
	Internasional 25	Nasional <input type="text"/>	
a. Kelengkapan Unsur Isi Buku (10%)	2.5		2.5
b. Ruang Lingkup & Kedalaman Pembahasan (30%)	7.5		7.5
c. Kecukupan & Kemutakhiran Data/Informasi & Metodologi (30%)	7.5		7.5
d. Kelengkapan Unsur & Kualitas Terbitan/Prosiding (30%)	7.5		7.5
<b>Total = (100%)</b>	<b>25</b>		<b>25</b>

**CATATAN PENILAIAN ARTIKEL OLEH REVIEWER:**

1. Tentang Kelengkapan dan Kesesuaian Unsur Isi Prosiding :

Artikel ini memiliki isi yg lengkap & sesuai

2. Tentang Ruang Lingkup dan Kedalaman Pembahasan :

Mung lingkup & pembahasan mendalam

3. Kecukupan & Kemutakhiran Data/Informasi & Metodologi :

Data & metodologi cukup jelas & baik

4. Kelengkapan Unsur & Kualitas Terbitan :

unsur & kualitas jurnal melalui prosiding ini cukup baik

Reviewer 2,



**Dr. Ir. Deden Sudrajat, M.Si**

NIK/NIDN : 0004096501

Jabatan Fungsional/Pangkat : Lektor Kepala, Pembina IV/a  
Unit Kerja : Universitas Djuanda

**LEMBAR**  
**REKAPITULASI PENILAIAN SEJAWAT SEBIDANG / PEER REVIEW**  
**KARYA ILMIAH : PROSIDING**

Judul Karya Ilmiah : **Production and Quality of Seven Basil (*Ocimum basilicum* L.) Accessions in Various Composition of Urea Fertilizer and Mexican Sunflower Compost**

Jumlah Penulis : 4 Orang

Status Pengusul : Penulis Pertama

Identitas Prosiding : a. Nama Prosiding : Proceedings of The International Seminar on Promoting Local Resources for Sustainable Agriculture and Development  
b. ISBN/ISSN : 978-94-6239-391-2/2468-7547  
c. Tahun Terbit, Lokasi : 11 Juni 2021  
d. Halaman/Penerbit : 267-273 / Atlantis Press  
e. Repository/Web : <https://www.atlantis-pers.com/proceedings/isplrsad-20/125957666>  
f. Terindeks di : CPCI, SCIE, ESCL, Google Scholar, Scopus

Kategori Publikasi Karya Ilmiah :  *Prosiding Forum Ilmiah Internasional*  
(beri ✓ pada kategori yang tepat)  *Prosiding Forum Ilmiah Nasional*

Hasil Penilaian *Peer Review* :

<b>Nilai Prosiding</b>		
<b>Peer Review 1</b>	<b>Peer Review 2</b>	<b>Nilai Rata-Rata</b>
<div style="border: 1px solid black; width: 40px; height: 40px; margin: auto; display: flex; align-items: center; justify-content: center;">25</div>	<div style="border: 1px solid black; width: 40px; height: 40px; margin: auto; display: flex; align-items: center; justify-content: center;">25</div>	<div style="border: 1px solid black; width: 40px; height: 40px; margin: auto; display: flex; align-items: center; justify-content: center;">25</div>
<p><b><u>KESIMPULAN :</u></b>  Nilai Karya Ilmiah Yang Diusulkan Ke Kopertis Wilayah IV Adalah : 25/</p>		



Part of **SPRINGER NATURE**

[PROCEEDINGS](#) | [JOURNALS](#) | [BOOKS](#)

---

Search

Series: [Advances in Biological Sciences Research](#)

## Proceedings of the International Seminar on Promoting Local Resources for Sustainable Agriculture and Development (ISPLRSAD 2020)

---

PROCEEDINGS OF THE INTERNATIONAL SEMINAR ON PROMOTING LOCAL RESOURCES FOR SUSTAINABLE AGRICULTURE AND DEVELOPMENT (ISPLRSAD 2020)

---

<

>

---

### Production and Quality of Seven Basil (*Ocimum basilicum* L.) Accessions in Various Composition of Urea Fertilizer and Mexican Sunflower Compost

#### Authors

Arifah Rahayu, Nur Rochman, Wini Nahraeni, Nida Fitriasaki

#### Corresponding Author

Arifah Rahayu

Available Online 11 June 2021.

#### DOI

<https://doi.org/10.2991/absr.k.210609.042> [How to use a DOI?](#)

#### Keywords

Basil, nitrate, vitamin C, chlorophyll, carotenoids

#### Abstract

The environmentally friendly fertilizer is necessary for vegetable production that are safe for consumption and maintain soil fertility. This study aims to determine the effect of the composition of urea fertilizer and Mexican

determine the effect of the composition of urea fertilizer and Mexican

sunflower compost (MSC) on the production and quality of various basil accessions (*Ocimum basilicum* L.). This study uses a factorial completely randomized design, namely basil accession (Gegerbitung, Kadudampit 1, Kadudampit 2, Kemang, Ciaruteun, Cijugug, and Gasol) and the composition of N fertilizer (100% N-urea, 100% N-MSC, 75% N-urea + 25% N-MSC, 50% N-urea + 50% N-MSC, 25% N-urea + 75% N-MSC, and 0% N-urea + 0% N-MSC). The

results showed that plants Kadudampit 2 and Gasol have growth and production better than other accessions. Plants treated with 100% N-MSC produced growth and harvest wet and dry weight not significantly different from 100% N-urea. Nitrate and vitamin C content were not significantly different between accessions. The use of 100% MSC significantly reduces leaf nitrate content. The vitamin C content was not significantly different between plants that were fertilized with various compositions of N. The content of chlorophyll a, chlorophyll b, and carotenoids of Gasol and Ciaruteun accessions were the lowest compared to the other five accessions. The highest chlorophyll a, chlorophyll b content are found in plants with Urea, followed by those given a combination of Urea + MSC. The carotenoid content of plant treated with 100% Urea was not significantly different from those fertilized with 75% Urea + 25% MSC and 25% Urea + 75% MSC.

#### Copyright

© 2021, the Authors. Published by Atlantis Press.

#### Open Access

This is an open access article distributed under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

[+ Download article \(PDF\)](#)



#### Volume Title

Proceedings of the International Seminar on Promoting Local Resources for Sustainable Agriculture and Development (ISPLRSAD 2020)

#### Series

Advances in Biological Sciences Research

**Publication Date**

11 June 2021

**ISBN**

978-94-6239-391-2

**ISSN**

2468-5747

**DOI**<https://doi.org/10.2991/absr.k.210609.042> [How to use a DOI?](#)**Copyright**

© 2021, the Authors. Published by Atlantis Press.

**Open Access**

This is an open access article distributed under the CC BY-NC license  
(<http://creativecommons.org/licenses/by-nc/4.0/>).

**Cite this article**

---

ris

enw

bib

---

TY - CONF  
AU - Arifah Rahayu  
AU - Nur Rochman  
AU - Wini Nahraeni  
AU - Nida Fitriasaki  
PY - 2021  
DA - 2021/06/11  
TI - Production and Quality of Seven Basil (*Ocimum basilicum* L.)  
Accessions in Various Composition of Urea Fertilizer and Mexican  
Sunflower Compost  
BT - Proceedings of the International Seminar on Promoting Local  
Resources for Sustainable Agriculture and Development (ISPLRSAD 2020)  
PB - Atlantis Press  
SP - 267  
FP - 272

LI 213

SN - 2468-5747

UR - <https://doi.org/10.2991/absr.k.210609.042>DO - <https://doi.org/10.2991/absr.k.210609.042>

ID - Rahayu2021

ER -

[+ download .ris](#)[COPY TO CLIPBOARD](#)

## Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: [contact@atlantis-press.com](mailto:contact@atlantis-press.com)

[▶ PROCEEDINGS](#)[▶ ABOUT](#)[▶ JOURNALS](#)[▶ NEWS](#)[▶ BOOKS](#)[▶ CONTACT](#)[▶ POLICIES](#)[▶ SEARCH](#)

---

[Home](#) [Privacy Policy](#) [Terms of use](#)

Copyright © 2006-2021 Atlantis Press – now part of Springer Nature

Part of **SPRINGER NATURE**

PROCEEDINGS | JOURNALS | BOOKS

---

Search

## Indexing Databases

---

### INDEXING DATABASES

---

Atlantis Press has affiliations with the following indexing databases:

---

**Web of Science**  
*Trust the difference*

### Web of Science

Clarivate Analytics' *Web of Science* is an online subscription-based citation indexing service which gives access to multiple databases that reference cross-disciplinary research and which allows for comprehensive citation search and in-depth exploration of specialized sub-fields within a scientific discipline. It consists of 6 core databases, a number of specialist collections, as well as regional databases and currently contains more than 160 million records and over 1.7 billion cited references. A select number of Atlantis Press journals and proceedings is indexed in Web of Science databases such as the *Science Citation Index Expanded* (SCIE), the *Emerging Sources Citation Index* (ESCI) and the *Conference Proceedings Citation Index* (CPCI).

---

**Scopus<sup>®</sup>** **Scopus**

Elsevier's *Scopus* is the world's largest abstract and citation database of peer-reviewed scientific journals, books and



conference proceedings which covers research topics across all scientific, technical and medical disciplines. The data

currently contains more than 75 million records and over 1.4 billion cited references, while it also offers various smart tools and metrics to track, analyze and visualize research. At present a select number of Atlantis Press journals and proceedings is indexed in Scopus and a number of applications are in progress.

---

## MEDLINE

### **MEDLINE**

*MEDLINE* is the **U.S. National Library of Medicine** premier bibliographic database that contains more than 25 million references to journal articles in life sciences with a concentration on biomedicine. The subject scope of MEDLINE is biomedicine and health, broadly defined to encompass those areas of the life sciences, behavioral sciences, chemical sciences and bioengineering needed by health professionals and others engaged in basic research and clinical care, public health, health policy development or related educational activities. MEDLINE also covers life sciences vital to biomedical practitioners, researchers and educators, including aspects of biology, environmental science, marine biology, plant and animal science as well as biophysics and chemistry. A select number of Atlantis Press journals in health and medical sciences is indexed in MEDLINE.



### **PubMed Central (PMC)**

*PubMed Central* (PMC) is a free full-text digital archive of biomedical and life sciences journal literature which has been developed and operated by the **National Center for Biotechnology Information** (NCBI), a division of the **U.S.**

**National Library of Medicine (NLM)** at the **U.S. National Institutes of Health (NIH)**. As of today, PMC contains more than 5.9 million full-text articles spanning several centuries of biomedical and life science research (late 1700s to present). Participation by publishers in PMC is voluntary, although participating journals must meet certain scientific and technical standards and content must be deposited as per the NIH Public Access Policy. A select number of Atlantis Press journals in health and medical sciences is indexed in PMC.

---



## Directory of Open Access Journals (DOAJ)

The *Directory of Open Access Journals* (DOAJ) is a community-curated online directory of open access journals which aims to be the starting point of all information searches for quality, peer-reviewed, open access material. DOAJ's mission is to increase the visibility, accessibility, reputation, usage and impact of quality, peer-reviewed, open access scholarly research journals globally, regardless of discipline, geography or language. At present, the directory contains more than 14,000 open access journals from 133 countries and more than 4.6 million open access articles covering all areas of science, technology, medicine, social science and humanities. All Atlantis Press journals are indexed in DOAJ.

---



## Ei Compendex

Elsevier's *Ei Compendex* (Engineering Index - COMPuterized ENgineering inDEX) on **Engineering Village** is the broadest and most complete engineering literature database in the world. It provides a holistic and global view of peer-reviewed and indexed publications with over 20 million records from 77 countries across 190 engineering disciplines. A select number

of Atlantis Press journals (i.e. within relevant subject areas) indexed in Ei Compendex.

---



## PubMed

*PubMed* is a free search engine which provides access to references and abstracts from biomedical and life sciences literature and which has been developed and maintained by the **National Center for Biotechnology Information** (NCBI), a division of the **U.S. National Library of Medicine** (NLM) at the **U.S. National Institutes of Health** (NIH). PubMed comprises more than 30 million citations and abstracts, primarily from **MEDLINE** but also from **PubMed Central** (PMC) and other life science journals and books, covering the fields of biomedicine and health and portions of the life sciences, behavioral sciences, chemical sciences and bioengineering. Many PubMed records also contain links to full-text articles, some of which are freely available (often in PMC). A select number of Atlantis Press journals in health and medical sciences is indexed in PubMed.

---



## Portico

*Portico* is a community-supported digital preservation archive that safeguards permanent access to e-journals, e-books and digital collections for researchers, scholars and students. The Portico digital preservation service is part of **ITHAKA**, a not-for-profit organization which aims to "help the academic community use digital technologies to preserve the scholarly record and to advance research and teaching in sustainable ways". The Portico archive currently contains more than 88 million journal articles, more than 1 million books and more than 4 million other digital collection items from a total number of 663 participating publishers. To

comply with NLM requirements (for indexing in **MEDL** and **PMC**) a select number of Atlantis Press journals is preserved in the Portico archive.

---



## China National Knowledge Infrastructure (CNKI)

The *China National Knowledge Infrastructure* (CNKI) is a key national information construction project launched by **Tsinghua University** and supported by the PRC Ministry of Education, PRC Ministry of Science and Technology, PRC Ministry of Propaganda and the PRC General Administration of Press and Publication. It is dedicated to the mass digitization of China knowledge resources as well as creating the platform for global dissemination and value-added services (CNKI was designated as the second agent of DOIs in mainland China by the **International DOI Foundation** in 2013). It comprises the China Integrated Knowledge Resources Database which contains over 90% of China knowledge resources. Most of the Atlantis Press proceedings and journals content is indexed in CNKI.

---



## Wanfang Data

*Wanfang Data* is an affiliate of the Chinese Ministry of Science & Technology which provides access to a wide range of database resources serving as a gateway to Chinese culture, medicine, business, science, engineering, etc. The Wanfang Data electronic resources comprise more than 43 million journal articles, close to 5 million dissertations and theses, more than 4 million conference papers and a broad collection of Chinese statistical data, patents, etc. Most of the Atlantis Press proceedings and journals content is indexed in Wanfang.



## dblp computer science bibliography

The *dblp computer science bibliography* is an online reference for bibliographic information on major computer science publications. Starting at the University of Trier in 1993 and owned by the Leibniz-Zentrum für Informatik since 2018, it evolved from a small experimental web server into a popular open-data service for the computer science community. The dblp mission is to support computer science researchers by providing free access to high-quality bibliographic metadata and links to electronic editions of publications. The dblp currently indexes more than 5.1 million publications, published by over 2.5 million authors. All Atlantis Press journals in fields related to computer science are indexed in the dblp.



## Ulrichsweb

ProQuest's *Ulrichsweb* is the standard online library directory and database for journals, magazines, newspapers and other periodicals. As such, it is regarded as the global authority for serials knowledge and analysis which comprises more than 383,000 serials from over 90,000 publishers covering 977 subject areas and 200 different languages. Records include searchable TOCs, ISSN, title, publisher, online availability, subject area, language, list prices and more. All Atlantis Press serial publications are indexed in Ulrichsweb.



## Google Scholar

*Google Scholar* is a freely accessible web search engine that indexes the full text or metadata of scholarly literature across an array of publishing formats and disciplines. The Google



Scholar index includes most peer-reviewed online academic journals and books, conference papers, theses and dissertations, preprints, abstracts, technical reports, and other scholarly literature, including court opinions and patents. It is estimated to contain more than 160 million documents and continues to cover approximately 90% of all articles published in English. All content published on the Atlantis Press platform is indexed in Google Scholar.

---

## Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: [contact@atlantis-press.com](mailto:contact@atlantis-press.com)

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ POLICIES
- ▶ ABOUT
- ▶ NEWS
- ▶ CONTACT
- ▶ SEARCH

---

[Home](#) [Privacy Policy](#) [Terms of use](#)



Copyright © 2006-2021 Atlantis Press – now part of Springer Nature

# Production and Quality of Seven Basil (*Ocimum basilicum* L.) Accessions in Various Composition of Urea Fertilizer and Mexican Sunflower Compost

Arifah Rahayu<sup>1\*</sup>, Nur Rochman<sup>1</sup>, Wini Nahraeni<sup>2</sup>, Nida Fitriasari<sup>1</sup>

<sup>1</sup>Agrotechnology Study Program, Faculty of Agriculture, University of Djuanda, Indonesia

<sup>2</sup>Agribusiness Study Program, Faculty of Agriculture, University of Juanda, Bogor, Indonesia

\* Corresponding author. Email: [arifah.rahayu@unida.ac.id](mailto:arifah.rahayu@unida.ac.id)

## ABSTRACT

The environmentally friendly fertilizer is necessary for vegetable production that are safe for consumption and maintain soil fertility. This study aims to determine the effect of the composition of urea fertilizer and Mexican sunflower compost (MSC) on the production and quality of various basil accessions (*Ocimum basilicum* L.). This study uses a factorial completely randomized design, namely basil accession (Gegerbitung, Kadudampit 1, Kadudampit 2, Kemang, Ciaruteun, Cijujung, and Gasol) and the composition of N fertilizer (100% N-urea, 100% N-MSC, 75% N-urea + 25% N-MSC, 50% N-urea + 50% N-MSC, 25% N-urea + 75% N-MSC, and 0% N-urea + 0% N-MSC). The results showed that plants Kadudampit 2 and Gasol have growth and production better than other accessions. Plants treated with 100% N-MSC produced growth and harvest wet and dry weight not significantly different from 100% N-urea. Nitrate and vitamin C content were not significantly different between accessions. The use of 100% MSC significantly reduces leaf nitrate content. The vitamin C content was not significantly different between plants that were fertilized with various compositions of N. The content of chlorophyll a, chlorophyll b, and carotenoids of Gasol and Ciaruteun accessions were the lowest compared to the other five accessions. The highest chlorophyll a, chlorophyll b content are found in plants with Urea, followed by those given a combination of Urea + MSC. The carotenoid content of plant treated with 100% Urea was not significantly different from those fertilized with 75% Urea + 25% MSC and 25% Urea + 75% MSC.

**Keywords:** Basil, nitrate, vitamin C, chlorophyll, carotenoids

## 1. INTRODUCTION

Basil (*Ocimum basilicum* L.) is a local vegetable that is commonly consumed fresh, mixed with cooking, and used as a biopharmaceutical or ornamental plant. Basil leaves have distinctive aroma from their essential oils. Traditionally, the essential oil of basil used as an antispasmodic, aromatic, carminative, digestive, and gastric agent [1]. The content of active compounds in basil leaves can optimize health and reduce the risk of chronic diseases such as coronary heart disease, cancer, and Alzheimer's [2].

Basil is an annual herbaceous plant (Blank *et al.* 2012) [3], bush-shaped. This plant grows upright until it reaches 100 cm in height [4]. The nutritional content in 100 g of basil leaves consists of 94 KJ of energy, 2.65 g of carbohydrates, 3.15 g of protein, 1.6 g of fiber, 0.64 g of fat, 0.8 mg of vitamin E, 18 mg of vitamin C, 5275 SI vitamin A, 177 mg Ca, 56 mg P, 3.17 mg Fe, 259 mg K,

0.81 mg Zn, 3142 µg beta-carotene, 0.034 mg thiamin, 0.092 mg niacin [5].

Growth, production, and quality of basil are influenced by climatic factors and cultivation techniques, including fertilization. A nutrient that plays an essential role in the growth of leafy vegetables is nitrogen. The application of nitrogen fertilizers can increase the production and fresh weight of basil [6]. Applying fertilizer with the right dose can increase plant height, dry weight, and essential oil content of basil plants [7]. However, too high an amount of nitrogen fertilizers can reduce plant height and leaf width, thus affecting wet weight and dried basil plants [6].

Plants absorbed nitrogen in the form of nitrate and ammonium. Generally, farmers use Urea as a nitrogen source, so that Urea is the most widely used nitrogen fertilizer globally, with a content of 46% N [8]. The application of Urea fertilizer can directly increase

plant productivity, especially leafy vegetables [9]. In rice, the dosage of Urea use by farmers can reach 400–600 kg/ha. Generally, farmers use urea as a nitrogen source, so that urea is classified as the most widely used nitrogen fertilizer in the world, with a content of 46% N [8]. The application of urea fertilizer can directly increase plant productivity, especially leafy vegetables [9]. In rice, the dosage of urea use by farmers can reach 400–600 kg/ha, exceeding the government's recommended dosage of 200-260 kg/ha [10].

The use of Urea fertilizers with high doses can increase the nitrate content in the soil, which causes groundwater pollution [11]. Too high absorption of nitrates by plants adversely affects the health of humans who consume them [12]. The use of environment from high chemical exposure can be done by providing a source of organic nitrogen, including using Mexican sunflower compost (MSC).

Mexican sunflower (*Tithonia diversifolia*) is a herbaceous plant from the Asteraceae family that used as green manure, liquid organic fertilizer, and compost [13]. Mexican sunflower contains 3.50 - 4.00% N, 0.35-0.38% P, 3.50 - 4.10% K, 0.59% Ca, and 0.27% Mg [14]. This plant has a faster decomposition time than others a high potential for restoring soil fertility [15]. Mexican sunflower compost can be an alternative to nitrogen fertilizers that can replace or reduce the use of Urea fertilizers. This study aims to evaluate the growth, production, and quality of various accessions of basil (*Ocimum basilicum* L.) on various composition of Urea and Mexican sunflower compost.

**2. MATERIALS AND METHODS**

The research was conducted from February to May 2020. The temperature conditions in the field ranged from 22 -34 °C, with an average humidity of 80%. The study took place during the rainy season, with an average rainfall of 517 mm/month [16]. The field experiment was carried out at the Experimental Garden of the Agrotechnology Study Program, Faculty of Agriculture, Djuanda University, Bogor. Analysis of leaf nitrate and vitamin C content was conducted in the laboratory of the Faculty of Agriculture, Djuanda University, leaf chlorophyll content in the Testing Laboratory of the Department of Agronomy and Horticulture of IPB, and analysis of nutrients and fertilizers at the ICBB (Indonesian Center for Biodiversity and Biotechnology) laboratory Bogor.

**2.1 Tools and Materials**

Materials used are basil accessions from Sukabumi district (Gegerbitung, Kadudampit1 and Kadudampit2), Bogor district (Kemang, Ciaruteun, and Cijujung), and Cianjur (Gasol), polybags, medium planting (soil and husk charcoal), MSC, synthetic fertilizers (Urea, SP- 36, and KCl), insecticides, distilled water, ingredients for analysis of vitamin C (starch and iodine solution), chlorophyll (extract *acetone* and Tris HCl 1%).

**2.2 Research Methods**

This study used a completely randomized design (CRD) consisting of two factors, namely basil accession and a combination of N fertilizers. Basil accessions consisted of seven levels, namely Gegerbitung, Kadudampit 1, Kadudampit II, Kemang, Ciaruteun, Cijujung, and Gasol. The combination of N fertilizer consists of six levels, namely 100% N-urea, 100% N-Mexican sunflower, 75% N-urea + 25% N-Mexican sunflower, 50% N-urea + 50% Mexican sunflower, 25% N-urea + 75% N-Mexican sunflower, 0% N-Urea + 0% N-Mexican sunflower. The dose of N fertilizer used was 150 kg N ha<sup>-1</sup>. The experiment consist of 42 treatment combinations, with three replications. Each experimental unit consists of four observation units, so there are 504 observation units (7 x 6 x 3 x 4).

Data were analyzed using variance (Test F). If the treatment had a significant effect, means compared with *Duncan's Multiple Range Test* (DMRT) at the 0.05 level.

**2.3 Research Implementation**

Mexican sunflower compost is done by mixing 25 kg of chopped fresh Mexican sunflower leaves with 100g of bran, 25g of granulated sugar dissolved in water, and 5ml of EM4. The mixed compost is stored in perforated *trash bag* and left for three weeks.

Sowing basil seeds for two weeks. Then the seeds are transferred to polybags measuring 10 × 20 cm. Transplanting the seedlings into polybags measuring 30 cm x 40 cm is done at the age of four weeks after seeding. The planting medium used for the nursery and planting is a mixture of soil and husk charcoal with a volume ratio of 1: 1. Watering, controlling pests, diseases, and weeds were carried out according to agricultural cultivation standards.

Fertilization is given based on the level of treatment. Urea fertilization is carried out gradually; 50% as a primary fertilizer and 25% as an additional fertilizer at the age of 3 and 6 weeks after planting (WAP). Mexican sunflower compost is given at once a week before planting, while SP-36 and KCl are given 100% as essential fertilizer.

Synthetic chemical fertilizer and MSC dosage per polybag were calculated by converting the fertilizer requirement per plant with a spacing of 50 x 25 cm (Table 1).

Table 1. Doses of chemical fertilizers and MSC for basil plants per polybag until the age of 8 WAP

Type of Fertilizer	Fertilizer dose	
	Per hectare	Per plant
Urea	326.09 kg/ha	5.43 g
SP-36	416.67 kg/ha	6.94 g
KCl	250.00 kg/ha	4.20 g
MSC	3846.2 kg/ha	64.1 g

Harvesting starts at the age of 4 weeks after planting (WAP) by cutting the young branches, 15 cm long. The harvest were carried out three times.

The variable observed were: plant growth (plant height, number of leaves, number of branches, the total length of branches, the diameter of stems, and leaf area). Leaf area was measured in the 5<sup>th</sup> widest leaf from the shoot at 6 WAP using the gravimetric method. Other variables observed were fresh weight and dry weight of harvest, fresh and dry weight of shoots and roots. The quality components observed were chlorophyll content by spectrophotometry, vitamin C content by iodine titration, and leaf nitrate content monitored by a nitrate meter (LAQUAtwin-NO3-11, HORIBA Scientific).

### 3. RESULTS AND DISCUSSION

#### 3.1 Results

##### 3.1.1 Soil Analysis and Organic Fertilizer

The results of soil analysis of medium planting showed that the soil used had a neutral pH with a value of 7.64, organic and total nitrogen contents were low (1.94%), while the C/N ratio is considered moderate (0.13). The available range of P<sub>2</sub>O<sub>5</sub> was 68.10 mg/g, P<sub>2</sub>O<sub>5</sub> potential, and K<sub>2</sub>O potential very high with values of 159.23 mg/100g and 117.53 mg/100g,

respectively. The soil criteria refer to Eviati and Sulaeman (2009) [17].

Analysis of organic fertilizers showed that Mexican sunflower compost contained N of 2.69%, P<sub>2</sub>O<sub>5</sub> 0.99%, and K<sub>2</sub>O 4.96%. The products are relatively low compared to the previous study [18] that N content of Mexican sunflower compost is 3.90%, and Lestari [14] that Mexican sunflower contains 3.50-4.00% N, 0.35- 0.38% P, 3.50-4.10% K. The total N content in compost has decreased because composting for 25-30 days with materials that have a high N content [19].

##### 3.1.2 Growth and Production of Basil Plants Basil

The Kadudampit 2 accession plant height was higher than the other accessions, but not significantly different from the Ciaruteun and Gasol accessions. Kemang accessions had no significant difference in leaf numbers with Gegerbitung, Kadudampit 2, and Cijujung, but significantly larger than other accessions. The number of shoots of ‘Kadudampit 1’ was higher than other accessions, but not significantly different from the ‘Gegerbitung’ and ‘Kemang’. The diameter of ‘Gasol’ was larger than the other accessions, but not significantly different from ‘Kadudampit2’ basil. The leaf area of ‘Gasol’ was larger than the other accessions but not significantly different from the Ciaruteun (Table 2).

**Table 2.** Plant height, number of leaves, number of shoots, the total length of shoots, stem diameter, and leaf area of basil plants

Treatment	Plant height (cm)	Number of leaves (strands)	Number of shoots (shoots)	The total length of shoots (cm)	Stem diameter (mm)	Leaf area (cm <sup>2</sup> )
<b>Accession</b>						
Gegerbitung	20:31 <sup>a</sup>	119.64 <sup>d</sup>	11.50 <sup>bc</sup>	95.27	4:45 <sup>a</sup>	22.92 <sup>a</sup>
Kadudampit 1	24.14 <sup>b</sup>	99.25 <sup>bc</sup>	12.42 <sup>c</sup>	105.55	5:03 <sup>ab</sup>	35.17 <sup>c</sup>
Kadudampit 2	29.47 <sup>c</sup>	113.93 <sup>cd</sup>	10.87 <sup>ab</sup>	103.15	5.73 <sup>bc</sup>	31.56 <sup>bc</sup>
Kemang	23:17 <sup>b</sup>	120.55 <sup>d</sup>	11.25 <sup>bc</sup>	93.05	4.64 <sup>a</sup>	27.85 <sup>ab</sup>
Ciaruteun	27.20 <sup>c</sup>	97.56 <sup>b</sup>	11.56 <sup>bc</sup>	93.31	5:27 <sup>ab</sup>	45.03 <sup>d</sup>
Cijujung	22.75 <sup>b</sup>	112.15 <sup>bcd</sup>	10.61 <sup>ab</sup>	98.12	4.85 <sup>ab</sup>	29.41 <sup>abc</sup>
Gasol	27.85 <sup>c</sup>	79.12 <sup>a</sup>	9.75 <sup>a</sup>	81.17	6:29 <sup>c</sup>	48.14 <sup>d</sup>
<b>Fertilizer Composition</b>						
100% Ua	27.36 <sup>b</sup>	113.96	11:49	104.95 <sup>bc</sup>	5:17	39.71 <sup>b</sup>
100% MSC	27.10 <sup>b</sup>	103.52	10:46	92.95 <sup>b</sup>	5:17	39.03 <sup>b</sup>
75% Ua+25% MSC	25.89 <sup>b</sup>	109.76	12:21	111.20 <sup>c</sup>	5.63	35.48 <sup>b</sup>
50% Ua+50% MSC	26.66 <sup>b</sup>	105.92	10.90	103.55 <sup>bc</sup>	5:06	35.85 <sup>b</sup>
25% Ua+75% MSC	25.58 <sup>b</sup>	108.54	10.67	94.14 <sup>b</sup>	4.93	34.60 <sup>b</sup>
0% Ua + 0% MSC	17.32 <sup>a</sup>	94.47	11.12	67.18 <sup>a</sup>	5.12	21.11 <sup>a</sup>

Means along the column with the same superscript are not significantly different by DMRT (p=0.05)

Ua: urea, MSC: Mexican sunflower compost

Plant height, the total length of shoots, and leaf area of basil plants given various compositions of Mexican sunflower fertilizers were not significantly different from those shown Urea 100%, but more significant than non fertilized N. The number of leaves, number of shoots, and stem diameter were not significantly different among the various N fertilizer compositions (Table 2).

The fresh and dry weight of Gasol accession was not significantly different from that of Kadudampit2. The fresh and dry weight of Kadudampit 2 accession roots The fresh weight of the ‘Kadudampit 2’ canopy was heavier than ‘Kemang’ and ‘Cijujung’, while the canopy dry weight did not differ between accessions (Table 3).

**Table 3.** Fresh and dry weight of harvesting, shoots, and roots of basil plants

Treatment	Harvesting weight (g)		Root weight (g)		Canopy weight (g)	
	Fresh	dry	fresh	Dry	Fresh	Dry
<b>Accession</b>						
Gegerbitung	47.02 <sup>ab</sup>	6.09 <sup>a</sup>	9.36 <sup>a</sup>	3.63 <sup>ab</sup>	131.60 <sup>bc</sup>	32.44
Kadudampit 1	51.45 <sup>bcd</sup>	6.37 <sup>a</sup>	6.70 <sup>a</sup>	2.78 <sup>a</sup>	114.54 <sup>abc</sup>	29.61
Kadudampit 2	55.94 <sup>de</sup>	7.46 <sup>b</sup>	13.40 <sup>b</sup>	4.82 <sup>b</sup>	168.12 <sup>c</sup>	40.24
Kemang	44.04 <sup>a</sup>	6.10 <sup>a</sup>	7.08 <sup>a</sup>	2.76 <sup>a</sup>	76.97 <sup>a</sup>	23.39
Ciaruteun	53.57 <sup>cd</sup>	6.68 <sup>ab</sup>	8.56 <sup>a</sup>	3.16 <sup>a</sup>	135.57 <sup>abc</sup>	31.16
Cijujung	48.79 <sup>abc</sup>	6.59 <sup>ab</sup>	7.71 <sup>a</sup>	3.08 <sup>a</sup>	101.25 <sup>ab</sup>	27.88
Gasol	60.21 <sup>e</sup>	7.38 <sup>b</sup>	8.69 <sup>a</sup>	3.31 <sup>a</sup>	125.38 <sup>abc</sup>	29.99
<b>Fertilizer composition</b>						
100% Ua	56.58 <sup>c</sup>	7.40 <sup>c</sup>	9.39	3.41 <sup>abc</sup>	117.69 <sup>b</sup>	30.52 <sup>bc</sup>
100% MSC	54.83 <sup>bc</sup>	7.16 <sup>bc</sup>	8.48	3.16 <sup>ab</sup>	118.31 <sup>b</sup>	26.02 <sup>b</sup>
75% Ua + 25% MSC	53.60 <sup>bc</sup>	6.95 <sup>bc</sup>	10.78	4.36 <sup>c</sup>	192.45 <sup>c</sup>	44.36 <sup>d</sup>
50% Ua + 50% MSC	51.37 <sup>b</sup>	6.80 <sup>bc</sup>	8.42	3.48 <sup>bc</sup>	122.33 <sup>b</sup>	34.53 <sup>cd</sup>
25% Ua + 75% MSC	50.78 <sup>b</sup>	6.51 <sup>b</sup>	8.74	3.21 <sup>ab</sup>	104.67 <sup>ab</sup>	30.83 <sup>bc</sup>
0% Ua + 0% MSC	42.30 <sup>a</sup>	5.18 <sup>a</sup>	7.18	2.54 <sup>a</sup>	76.05 <sup>a</sup>	17.76 <sup>a</sup>

Means along the column with the same superscript are not significantly different by DMRT (p=0.05)

Ua: urea, MSC: Mexican sunflower compost

The fresh and dry weight of harvest, roots, and canopy of plants shown 100% MSC was not significantly different from those given 100% Urea, as well as those given composition of 75% Urea + 25% MSC, even fresh and dry weight. The canopy of plants given this fertilizer composition more significant than those shown 100% Urea.

### 3.1.3 Quality of Basil Leaves

The quality of basil leaves (content of nitrate and vitamin C) were significantly influenced by fertilizer's composition. Plants treated with 100% Urea showed the highest nitrate content, not substantially different from those given 25% Urea + 75% MSC, in comparison, plants treated with 100% MSC and not N fertilization showed the lowest nitrate content (Table 5). The content of vitamin C in N fertilizers' various compositions was not significantly different, but substantially higher than plant without N fertilizer (Table 4).

The pigment content of chlorophyll a, chlorophyll b, carotene, and total chlorophyll was significantly affected by accession treatment and fertilizer composition, while anthocyanin content was not affected. Basil leaves of 'Kemang' contain chlorophyll a, chlorophyll b, total chlorophyll, and carotene higher compared to Gasol and Kadudampit1. Basil plants treated with 100% Urea fertilizer composition show the highest content of chlorophyll a, chlorophyll b, total chlorophyll, and carotene, followed by the composition of a mixture of Urea + MSC, while the plants treated with 100% MSC

and those not given N fertilizers had the lowest pigment content (Table 5).

**Table 4.** The content of nitrates and vitamin C of basil leaves

Treatment of	Nitrate	Vitamin C
<b>Accession</b>		
Gegerbitung	94.52	1818.50
Kadudampit 1	1755.50	94.52
Kadudampit 2	1883.30	88.30
Kemang	92.74	1740.70
Ciaruteun	2079.60	87.70
Cijujung	96.30	1649.20
Gasol	99.56	1888.80
<b>Fertilizer Composition</b>		
100% Ua	2373.0 <sup>c</sup>	99.80 <sup>b</sup>
100% MSC	1358.0 <sup>a</sup>	96.76 <sup>b</sup>
75% Ua + 25% MSC	1957.1 <sup>b</sup>	97.02 <sup>b</sup>
50% Ua + 50% MSC	1888.8 <sup>b</sup>	94.73 <sup>b</sup>
25% Ua + 75% MSC	2073.0 <sup>bc</sup>	94.98 <sup>b</sup>
0% Ua + 0% MSC	1334.9 <sup>a</sup>	76.95 <sup>a</sup>

Means along the column with the same superscript are not significantly different by DMRT (p=0.05)

Ua: urea, MSC: Mexican sunflower compost



**Table 5.** Content Pigment of basil plants

Treatment	Leaf pigment content (mg.g <sup>-1</sup> )				
	Chlorophyll a	Chlorophyll b	Anthocyanin	Carotenoids	Chlorophyll Total
<b>Accession</b>					
Gegerbitung	1.62 <sup>cd</sup>	0.84 <sup>c</sup>	0.04	0.48 <sup>bc</sup>	2.45 <sup>c</sup>
Kadudampit I	1.51 <sup>bc</sup>	0.77 <sup>bc</sup>	0.05	0.42 <sup>ab</sup>	2.28 <sup>bc</sup>
Kadudampit II	1.67 <sup>cd</sup>	0.83 <sup>c</sup>	0.02	0.48 <sup>bc</sup>	2.50 <sup>c</sup>
Kemang	1.75 <sup>d</sup>	0.87 <sup>c</sup>	0.04	0.49 <sup>c</sup>	2.62 <sup>c</sup>
Ciaruteun	1.33 <sup>ab</sup>	0.68 <sup>ab</sup>	0.04	0.40 <sup>a</sup>	2.01 <sup>ab</sup>
Cijujung	1.63 <sup>cd</sup>	0.83 <sup>c</sup>	0.08	0.43 <sup>abc</sup>	2.46 <sup>c</sup>
Gasol	1.27 <sup>a</sup>	0.63 <sup>a</sup>	0.01	0.38 <sup>a</sup>	1.90 <sup>a</sup>
<b>Fertilizer composition</b>					
100% Ua	1.92 <sup>c</sup>	0.95 <sup>c</sup>	0.04	0.52 <sup>c</sup>	2.88 <sup>c</sup>
100% MSC	1.23 <sup>a</sup>	0.63 <sup>a</sup>	0.08	0.35 <sup>a</sup>	1.88 <sup>a</sup>
75% Ua + 25% MSC	1.62 <sup>b</sup>	0.84 <sup>b</sup>	0.05	0.48 <sup>bc</sup>	2.46 <sup>b</sup>
50% Ua + 50% MSC	1.55 <sup>b</sup>	0.78 <sup>b</sup>	0.03	0.44 <sup>b</sup>	2.33 <sup>b</sup>
25% Ua + 75% MSC	1.62 <sup>b</sup>	0.83 <sup>b</sup>	0.02	0.48 <sup>bc</sup>	2.45 <sup>b</sup>
0% Ua + 0% MSC	1.27 <sup>a</sup>	0.64 <sup>a</sup>	0.02	0.37 <sup>a</sup>	1.91 <sup>a</sup>

Means along the column with the same superscript are not significantly different by DMRT (p=0.05)

### 3.2 DISCUSSION

#### 3.2.1 Growth and Production of Basil Plants

Basil accessions in this study can be distinguished on small leaves (Gegerbitung, Kemang, and Cijujung) and broad leaves (Kadudampit1, Kadudampit2 Ciaruteun, and Gasol). Accessions of broad-leaves basil had significantly higher yields on the variables of plant height, stem diameter, and leaf area. Still, they had the lower number of leaves, the number of the shoot, and the total length of shoots compared to accessions with small leaves, except for Kadudampit2 accession because Kadudampit2 accession began to appear leafy width at 4 WAP. This related to translocation of photosynthate to the part of plant growth [20]. The photosynthate of broadleaf basil accession is allocated more to leaf size than to the number of leaves, the number of shoots, and total length of shoots.

Basil is a leafy vegetable, so the number of leaves, the number and the length of shoots affect the productivity of the basil plant. The ‘Gasol’ harvest fresh weight was significantly higher than other accessions, because ‘Gasol’ had the largest leaf size compared to different accessions. Plant with broad leaves will be able to absorb more sunlight, thus increasing photosynthetic rate, and affecting plant growth [21]. This following Gigir *et al.* (2014) [22], which states that the taller the plant, the longer and broader the leaves, the greater the yield.

Plants treated with N fertilizers showed better growth and production responses than those not fertilized. The results of Damayanti’s (2017) [23] research show that nitrogen application has a significant effect on the height of the basil plant. According to Aziz and Kurnia (2015), sufficient nitrogen supply to plants can accelerate plant vegetative growth, both on stems, branches, and leaves. The growth and production of basil plants treated of MSC or combined with Urea were not significantly different. This results showed that MSC can substitute or complement

for Urea. This related to the ability of MSC to improve soil physical and chemical properties. Mexican sunflower green manure can change the physical properties (reduce the soil density) and chemical properties (increase the content of C-organic, N-total, available-P and K interchangeable) soil, and increase the fresh weight of cabbage flowers [24].

The provision of inorganic fertilizers combined with organic fertilizers can respond to increased growth of basil plants [22]. Sulaeman *et al.* [25] also showed that the application of organic fertilizers (manure or sludge) combined with synthetic fertilizers resulted in higher plant growth and higher maize plant residues. Kurnia *et al.* [26] reported that the production of tomato plants given 30 ton.ha<sup>-1</sup> grass compost and 15 ton.ha<sup>-1</sup> grass compost + 50% synthetic fertilizer dose was higher than control.

Basil plants treated without N fertilizer (0% Ua + 0% MSC) showed the lowest growth and production compared to other treatments. This related to the low nitrogen content of the soil used for research, namely 0.13%, so it requires additional elements in the form of fertilizer.

#### 3.2.2 Quality of Treated Basil Plants

Plant leaves that fertilized by 100% Urea, showed the highest nitrate content. This following reported of Liu *et al.* [27] that lettuce treated with inorganic fertilizers had a higher nitrate concentration reaching 5000-6100 mg/kg, when compared to organic fertilizer lettuce, namely 4300-5200 mg/kg. This because the Urea used will immediately hydrolyze and undergo a nitrification process to form NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> ions .

Nitrate and nitrite levels in vegetables can vary due to the use of fertilizers and biotic, and abiotic factors. Continuous use of inorganic fertilizers will accumulate in the soil so that nitrate levels will increase and vegetables

tend to have high nitrate levels. . These condition needs to be watched out because the estimated total average intake of nitrate consumed is around 61 mg/person/day (range 24-102 mg/person/day), and vegetables contribute around 75% [28].

The application of MSC makes the nitrate content in basil leaves low. This one of the advantages of MSC because vegetables with low nitrate levels are safer for human consumption. Plants treated with various composition of Urea show a higher content of vitamin C compared to non-fertilized. Presumably this is caused by nitrogen from the composition of the fertilizers relatively sufficient for formation of leaf chlorophyll, thus contributing to increasing the rate of photosynthesis. Plants used photosynthate in the form of glucose (C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>) as a precursor for the formation of vitamin C and other chemical compounds [29].

The mainstay of treatment compared to other treatments. Presumably this condition related to time offering MSC, which is only done once a week before planting, while Urea is given gradually as essential fertilizer, at the age of 3 and 6 weeks after planting.

The total chlorophyll a and b content in this study were relatively low, about 1.90 - 2.88. Basil has a low range because it is an annual plant. The leaves are small with thin leaves. This is following to Fahn (1982) [30], which states that leaves basil are only composed of one palisade tissue layers. Mesophyll, especially palisade tissue, is a network that is rich in chlorophyll.

Chlorophyll and carotene pigments are located in chloroplasts and play a role in the photosynthesis process. Carotene can protect chlorophyll from high light intensity [31]. The results showed that there was a link between carotene and chlorophyll. Raisawati (2018) [32] showed a correlation between the two; an increase in carotene levels related to an increase in chlorophyll levels.

#### 4. CONCLUSION

Kadudampit2 and Gasol accessions can be developed because they show plant height, fresh weight, and dry weight greater than other accessions. The application of MSC and its combination with Urea showed growth, production and quality equivalent to those treated with Urea. MSC can reduce the nitrate levels absorbed by plants, making it safer to consumed and can improve soil fertility. The treatment of 75% N-Urea and 25% N- Mexican sunflower can be an alternative fertilizer that can be used for basil plants because it can balance several aspects both in terms of productivity, economy, and environment.

#### ACKNOWLEDGMENTS

Thank you to the Directorate General of Higher Education, Ministry of Education and Culture of the Republic of Indonesia for research funding assistance through the 2019 Higher Education Excellence Applied Research Grant.

#### REFERENCES

- [1] S. Khan *et al.*, "Phytochemical Constituents and Pharmacological Activities of Sweet Basil- *Ocimum basilicum* L. ( Lamiaceae )," vol. 23, no. 9, pp. 3773–3782, 2011.
- [2] A. Guaadaoui, S. Benaicha, and N. Elmajdoub, "What is a bioactive compound? A combined definition for a preliminary consensus," no. April, 2015.
- [3] A. F. Blank *et al.*, "A diallel study of yield components and essential oil constituents in basil (*Ocimum basilicum* L.)," *Ind. Crops Prod.*, vol. 38, pp. 93–98, 2012.
- [4] D. N. Nuraini, *Aneka daun berkhasiat untuk obat*. Yogyakarta: Gava Media, 2014.
- [5] U. S. D. A. [USDA], "Ocimum basilicum L.," 2018. [Online]. Available: <https://fdc.nal.usda.gov/fdc-app.html#/food-details/172232/nutrients>. [Accessed: 29-Apr-2020].
- [6] R. N. Wierdak, E. Rozek, and K. Dzida, "Growth response to nitrogen and potassium fertilization of common basil (*Ocimum basilicum* L.) Plants," vol. 11, no. 2, pp. 275–288, 2012.
- [7] K. Singh, P. P. Singh, S. U. Beg, D. Kumar, and D. D. Patra, "Effect of NPK fertilizers on growth, oil yield and quality of French basil (*Ocimum basilicum* L)," *J. Aromatic and Spices Crops*, vol. 13, no. 1, pp. 52–54, 2004.
- [8] A. Barker and D. Pilbeam, *Nitrogen*. In Barker AV, Pilbeam DJ (Eds.). *Handbook of Plant Nutrition*. Boca Raton: CRC Press. Taylor and Francis Group., 2007.
- [9] Endrizal and J. Bobihoe, "Efisiensi penggunaan pupuk nitrogen dengan penggunaan pupuk organik pada tanaman padi sawah," *J. Pengkajian dan Pengembangan Teknologi Pertanian*, vol. 7, no. 2, pp. 118–124, 2004.
- [10] A. S. Wahid, "Peningkatan efisiensi pupuk nitrogen pada padi sawah dengan metode bagan warna daun," *J. Penelitian dan Pengembangan Pertanian*, vol. 22, no. 1999, 2003.
- [11] L. M. Purba, T. Ashar, and D. N. Santi, "Pola Pemupukan dan Analisa Kandungan Nitrat pada Sayur Brokoli (*Brassica oleracea* cv. *brocoli*) di Pertanian Desa Merdeka Kecamatan Merdeka Kabupaten Karo Tahun 2012," pp. 1–9, 2012.
- [12] W. Safitri, R. S. Pujiati, and P. T. Ningrum, "Kandungan nitrat pada air tanah di sekitar lahan pertanian padi, palawija dan tembakau," *Artikel ilmiah Hasil Penelitian Mahasiswa*, 2014.
- [13] M. Nurrohman, A. Suryanto, and K. Puji, "Penggunaan Fermentasi Ekstrak Paitan (*Tithonia diversifolia* L.) dan Kotoran Kelinci Cair Sebagai Sumber Hara Pada Budidaya Sawi (*Brassica juncea* L.) Secara Hidroponik Rakit Apung," *J. Produksi Tanam.*, vol. 2, no. 8, pp. 649–657, 2014.
- [14] S. A. D. Lestari, M. Melati, and H. Purnamawati, "Penentuan Dosis Optimum Pemupukan N, P, dan K

- pada Tanaman Kacang Bogor [*Vigna subterranea* (L.) Verdcourt],” *J. Agron. Indones. (Indonesian J. Agron.)*, vol. 43, no. 3, p. 193, 2016.
- [15] T. S. Rahardian, T. Sumarni, and A. Suryanto, “Pemanfaatan pupuk hijau paitan (*Tithonia diversifolia*) dan krinyu (*Chromolaena odorata*) dalam peningkatan hasil tanaman brokoli (*Brassica oleracea*)” *Planta Tropica* vol. 2, no. 2, pp. 108–116, 2017.
- [16] BMKG, *Analisis Hujan Februari 2020 dan Prakiraan Hujan April, Mei dan Juni 2020*. Jakarta, 2020.
- [17] Eviati and Sulaeman, *Analisis Kimia Tanah, Tanaman, Air, dan Pupuk*. Bogor: Balai Penelitian Tanah, 2009.
- [18] A. Rahayu, W. Nahraeni, N. Rochman, and A. Faturrochman, “respon pertumbuhan aksesi kemangi pada berbagai komposisi pupuk nitrogen alami” *J. Agronida*, vol. 5, no.2, pp 70-77, 2019.
- [19] A. Arifiati, Syekhfani, and Y. Nuraini, “Uji Efektifitas Perbandingan Bahan Kompos Paitan (*Tithonia Diversifolia*) Tumbuhan Paku (*Dryopteris Filixmas*) dan Kotoran Kambing Terhadap Serapan N Tanaman Jagung Pada Inceptisol,” *J. Tanah dan Sumberd. Lahan*, vol. 4, no. 2, pp. 543–552, 2017.
- [20] Hermansyah, Y. Sasmita, and E. Inorah, “Penggunaan pupuk daun dan manipulasi jumlah cabang yang ditinggalkan pada panen kedua tanaman nilam,” *Akta Agrosia*, vol. 12, pp. 194–203, 2009.
- [21] H. B. Buntoro, R. Rogomulyo, and S. Trisnowati, “Pengaruh takaran pupuk kandang dan intensitas cahaya terhadap pertumbuhan dan hasil temu putih (*Curcuma zedoaria* L.)” *Vegetalika*, vol. 3, no. 4, pp. 29–39, 2014.
- [22] S.F. Gigir, J. J. Rondonuwu, W. J. N. Kumolontang, dan Rafli I. Kawulusan, “Respons pertumbuhan kemangi (*Ocimum sanctum* L.) terhadap pemberian pupuk organik dan anorganik,” *Cocos*, vol. 5, no. 3, pp 2014.
- [23] D. P. O. Damayanti, “Pengaruh ammonium (NH<sub>4</sub><sup>+</sup>) dan nitrat (NO<sub>3</sub><sup>-</sup>) terhadap pertumbuhan dan kandungan minyak atsiri tanaman kemangi (*Ocimum basilicum*) dengan sistem hidroponik,” Unoversitas Jember, 2017.
- [24] Hafifah, Sudiarmo, M. . Maghfoer, and B. Prasetya, “The potential of *Tithonia diversifolia* green manure for improving soil quality for cauliflower (*Brassica oleracea* var. *botrytis* L.)” *Journal Degrad. Min. Lands Manag.*, vol. 3, no. 2, pp. 499–506, 2016.
- [25] Y. Sulaeman, Maswar, and D. Erfandi, “Effect of a combination of organic and inorganic fertilizers on soil chemical properties and yield of maize on acid dry land,” *J. Assess. Tech Dev. Pert.*, vol. 20, no. 1, pp. 1–12, 2017.
- [26] S. Kurnia, N. Setyowati, and Alnopri, “The effect of combined doses of weed compost and synthetic fertilizers on the growth and yield of tomato (*Lycopersicon esculentum* Mill.)” *J. Pert Sci.*, vol. 21, no. 1, pp. 15–21, 2019.
- [27] C.W. Liu, Y. Sung, B.C. Chen, H.Y. Lai, “Effects of nitrogen fertilizers on the growth and nitrate content of lettuce (*Lactuca sativa* L.)” *Int. J. Environment. Res. and Public Health*, vol. 11, no. 4, pp. 4427–4440, 2014.
- [28] J. M. Hill, *Nitrates and Nitrites in Food and Water*. Cambridge: Woodhead Publishing Limited, 1996.
- [29] R. Novitasari, “Proses Respirasi Seluler Pada Tumbuhan,” *Pros. Semin. Nas. Pendidik. Biol. dan Biol.*, pp. 89–96, 2017.
- [30] A. Fahn, *Anatomi Tumbuhan*, vol. 3, no. 1. Semarang: Gadjah Mada University Press, 1982.
- [31] C. Pebrianti, R. B. Ainurrasyid, L. Purnamaningsih, R. Leaf, and B. Merah, “Uji kadar antosianin dan hasil enam varietas tanaman bayam merah (*Alternanthera amoena* Voss.) Pada Musim Hujan,” *J. Produksi Tanam.*, vol. 3, no. 1, pp. 27–33, 2015.
- [32] T. Raisawati, “Kadar total klorofil, karoten, antosianin dan vitamin C tempuyung (*Sonchus arvensis* L.) pada cara panen berbeda,” *Prosiding Seminar Nasional LPPM UNiversitas Negeri Surabaya*, July, pp. 729–736, 2018.