

External and Internal Quality of the Quail Eggs Fed Ration with Different Level of Metabolizable Energy and Protein

By Dede Kardaya

External and Internal Quality of the Quail Eggs Fed Ration with Different Level of Metabolizable Energy and Protein

Deden Sudrajat*, Dewi Wahyuni, Hanafi Nur, Dede Kardaya
Department of Animal Science, Faculty of Agriculture University of Djuanda
Jl. Tol Ciawi 1 Kotak pos 35 Bogor, Indonesia

*Corresponding author: deden.sudrajat@unida.ac.id

(Submitted: November 19, 2021; Accepted: February 05, 2022)

ABSTRACT

Quail eggs are a source of animal protein that the public can consume. In addition to the low price, eggs also contain nutrients needed by the body. So it is necessary to pay attention to the quality of the eggs before consuming them. Egg quality that can be assessed is derived from internal and external quality. The quality of the eggs is influenced by various factors, including the content of the ratio given, especially the energy-protein balance of the feed. This study aimed to assess the external and internal quality of quail with different protein and energy levels. The research was carried out for 42 days in the poultry house of the Animal Husbandry Study Program, Faculty of Agriculture, Djuanda University. The feed ingredients formulated in this study were yellow corn, fine bran, soybean meal, fish meal, premix, DCP, CaCO₃, CPO, and Cr-O. The design used was a completely randomized design (CRD) with four treatments and four replications with six quails per cage unit. The research data were analyzed by analysis of variance (ANOVA) if the data showed significantly different results ($P < 0.05$), followed by Duncan's test. The variables observed in this study were egg weight, shell weight, percentage of eggshell weight, eggshell thickness, egg index, egg white weight, egg yolk weight, egg white weight percentage, egg yolk weight percentage, Haugh unit (HU). The results showed that the balance of energy and protein differences were not significantly different in all variables. Based on the research results, it can be concluded that the quail ration with the balance of protein 17% and metabolizable energy 2800 kcal/kg has been able to maintain the internal and external quality of eggs in quail and produce standard egg quality.

Keywords: protein energy balance; egg white, yolk, haugh unit

INTRODUCTION

Quail is one of the poultry that can produce eggs and meat. Quail is poultry that can be developed easily because quail is very easy to reared and can grow and develop quickly (Lari et al. 2016). One type of quail can be kept in the Japanese quail (*Coturnix coturnix japonica*). Female quail can be used as seeds and quail egg producers. Laying quail for consumption are eggs that are not fertilized and produced by females. On the contrary, quail for quail-brooders are eggs that can be incubated and the eggs produced are fertilized (Panekanan et al., 2013). The

national quail population continues to increase in 2019 to 14,107,479 quails (Ditjen PKH, 2019).

Eggs are a source of animal protein that the public can consume. In addition to the low price, eggs also contain nutrients needed by the body. So it is necessary to pay attention to the quality of the eggs before consuming them. Egg quality that can be assessed is derived from eggs' internal and external quality. External quality can be seen from the thickness of the shell, egg index, egg weight, while the internal quality can be seen from the percentage of egg white weight, egg yolk percentage, Haugh unit. The quality of the eggs is influenced by various factors,



including the content of the ration given at the time of rearing the quail. Besides affecting egg quality and production, feed is the most crucial in production costs because 60-80% of expenses are used as feed costs (Khalil 2015). This causes farmers to face difficulties in providing quality feed that can meet the nutritional needs of quail. It can be overcome this problem by determining the standard of quail nutrient needs, especially the energy-protein balance of feed.

Energy-protein balance is the most critical factor affecting feed quality and determining quail productivity (Napirah et al., 2018). Although the energy content of the feed is required to be balanced with the protein content of the feed, low protein feeds can decrease body weight (Kamran et al., 2008). In contrast, high protein feeds can increase quail body weight gain and internal quality in quail eggs (Kamran et al., 2008). Gheisari et al., 2011). Therefore, research on the provision of different energy and protein balances needs to be carried out in this study to improve quail eggs' external and internal quality. This study aims to assess the external and internal quality of quail eggs with different protein and energy levels.

MATERIAL AND METHOD

This research was carried out for 42 days in the Poultry house of the Department of Animal Science, Faculty of Agriculture, Djuanda University, Bogor. The tools and materials used in this study were 96 female quails aged 48 days with an initial weight of 151.43±6.81 g. The feed provided was self-mixing

consisting of yellow corn, fine bran, soybean meal, fish meal, premix, DCP, CaCO₃, CPO, and Cr-O. The composition of the feed ingredients used as quail rations in this study can be seen in Table 1. The cage used was a battery cage where there were 16 cage plots.

The design used was a completely randomized design with four treatments, four replications, and each experimental unit consisted of 6 quails. The treatments carried out in this study were as follows: R1 = Crude Protein 17%, Metabolizable Energy 2800 kcal/kg. R2 = Crude Protein 18%, Metabolizable Energy 2950 kcal/kg. R3 = Crude Protein 19%, Metabolizable Energy 3000 kcal/kg. R4 = Crude Protein 20%, Metabolizable Energy 2900 kcal/kg. Data were analyzed using ANOVA with the help of the SPSS 25 application and further tested using Duncan Multiple test.

The variables in this study were the external quality of eggs (egg weight, shell weight, shell thickness, and egg index) and internal egg quality (egg white weight percentage, egg yolk weight percentage, and Haugh units). The cages used for research were cleaned with a disinfectant solution. Feeding of 20 g/head/day given two times a day at 08.00 and 16.00, providing drinking water, cleaning of excreta, and sanitation around the cage. After that data collection, data was taken once a week for three weeks, carried out in the 3rd study week from the beginning of the study with 62 days of quail age until the fifth week of the study.

Table 1. Composition and nutrient content of experimental ration

| No | Ingredients | Amount % | | | |
|---------|--------------------------------|----------|------|------|------|
| | | R1 | R2 | R3 | R4 |
| 1 | Yellow corn | 63 | 63.5 | 62.5 | 62.0 |
| 2 | Rice Bran | 5,5 | - | - | - |
| 3 | Soybean Meal | 17.0 | 19.0 | 15.9 | 17.4 |
| 4 | Fish meal | 5,4 | 6.4 | 10 | 11.0 |
| 5 | Premix | 1.0 | 1.0 | 1.0 | 1.0 |
| 6 | CaCO ₃ | 5.0 | 5.0 | 5.0 | 5.0 |
| 7 | CPO | 0,5 | 2.5 | 3.0 | 1.5 |
| 8 | DCP | 2,5 | 2.5 | 2.5 | 2.0 |
| 9. | Cr-O | 0,1 | 0.1 | 0.1 | 0.1 |
| | Total | 100 | 100 | 100 | 100 |
| Nutrien | | | | | |
| | Crude Protein (%) | 17 | 18 | 19 | 20 |
| | Metabolizable Energy (Kkal/kg) | 2800 | 2950 | 3000 | 2900 |

Note: CaCO₃ (lime), CPO = Crude Palm Oil (oil), DCP = Dicalcium Phosphate, Cr-O = Organic Chromium.

22 RESULT AND DISCUSSION

The effect of energy and protein balance on quail eggs' external and internal quality can be seen in Tables 2 and 3. The results were not significantly different ($p>0.05$) on all variables.

There are three indicators to determining quail eggs' external quality: egg weight, egg index, and eggshell thickness. The weight of the eggs is essential to be considered because the heavier the eggs, the eggs tend to have a high economic value and are more attractive to consumers. A good egg index is needed for egg grading. At the same time, a suitable shell thickness will minimize the inside of the egg being contaminated with the environment so that the egg's contents have good quality, and stay awake (Nugraha et al., 2018).

The analysis of variance showed that the results were not significantly different ($P>0.05$) in egg weight. The average egg weight in this study was 8.99-9.99 g. This study is lower than Ardianyah et al. (2016), which provides different protein levels, namely 9.76-11.56 g. According to Parizidian et al. (2011), the egg weight ranged from 10-11.9 g or about 8% of the bodyweight of the parent. In the ratio, egg weight is influenced by genetics and protein

(Sujana et al., 2020). The protein used differed from 17 to 20% of the dose in this study but did not give significantly different results. Therefore, it is suspected that the protein requirement of 17% has been met. Quail at laying eggs requires energy and protein for bare life, egg production, and growth. After the basic needs of life are met, the energy and protein consumed tend to be used for production rather than increasing egg weight (Sujana et al. 2020).

The shell is the outer egg structure that reduces physical and biological damage to the egg (Sujana et al. 2020). The average shell weight in this study ranged from 0.89-0.92 g. The thickness of the shell is one of the factors that affect egg quality because the shell can protect the contents of the egg. In addition, the thickness of the shell is much influenced by the level of calcium in the ratio, which determines the availability of calcium salts in the blood for egg formation (Yuwanta 2010). The average thickness of the shell in this study was 0.18-0.19 mm.

Means egg index was in the range of 78.10-79.40%. For comparison, research by Nugraha et al. (2018) produced an egg index of 77.62-78% in quail fed a ration containing 22% protein.

Table 2. Means external quality of eggs

| treatment | Variabels | | | |
|-----------|----------------|----------------|------------------|----------------------|
| | Egg Indeks (%) | Egg Weight (g) | Shell Weight (g) | Shell Thickness (mm) |
| R1 | 78,25 ± 1,45 | 8,99 ± 0,31 | 0,89 ± 0,02 | 0,18±0,008 |
| R2 | 78,45 ± 1,36 | 9,21 ± 0,54 | 0,87 ± 0,35 | 0,18±0,007 |
| R3 | 78,10 ± 1,21 | 9,29 ± 0,34 | 0,91 ± 0,45 | 0,18±0,008 |
| R4 | 79,40 ± 0,65 | 9,99 ± 0,60 | 0,92 ± 0,40 | 0,19±0,003 |

Note: R1 = Crude Protein 17%, Metabolizable Energy 2800 kcal/kg, R2 = Crude Protein 18%, Metabolizable Energy 2950 kcal/kg, R3 = Crude Protein 19%, Metabolizable Energy 3000 kcal/kg, R4 = Crude Protein 20%, Metabolizable Energy 2900 kcal/kg.

Table 3. Means internal quality of quail eggs

| Variabels | Treatments | | | | |
|----------------------|--------------|--------------|--------------|--------------|--------------|
| | R1 | R2 | R3 | R4 | Means |
| Egg white weight (g) | 5.09 ± 0.23 | 4.93 ± 0.59 | 5.00 ± 0.35 | 5.63 ± 0.44 | 5.16 ± 0.47 |
| Egg yolk (g) | 2.71 ± 0.15 | 2.65 ± 0.32 | 2.57 ± 0.30 | 2.81 ± 0.28 | 2.69 ± 0.26 |
| Egg white weight (%) | 55.39 ± 0.64 | 54.54 ± 1.38 | 58.37 ± 6.05 | 57.07 ± 1.30 | 56.34 ± 3.24 |
| Egg yolk (%) | 30.32 ± 1.20 | 29.17 ± 0.96 | 28.61 ± 1.39 | 28.31 ± 1.60 | 29.10 ± 1.41 |
| Haugh Unit | 82.11 ± 1.39 | 84.04 ± 1.59 | 84.41 ± 0.52 | 84.83 ± 1.91 | 83.85 ± 1.68 |

Notes: R1 = Crude Protein 17%, Metabolizable Energy 2800 kcal/kg, R2 = Crude Protein 18%, Metabolizable Energy 2950 kcal/kg, R3 = Crude Protein 19%, Metabolizable Energy 3000 kcal/kg, R4 = Crude Protein 20%, Metabolizable Energy 2900 kcal/kg.

The statistical analysis results showed that the treatment was not significantly different ($P>0.05$) on egg white weight, egg yolk weight, egg white weight percentage, egg yolk weight percentage, and Haugh unit. These were thought to be caused because the protein content in the ration given to quail has met the need to achieve optimal egg weight (Agustini et al., 2014). According to Wahju (2004), egg quality is influenced by various factors, including genetic and environmental factors, nutrient content, environmental temperature, disease, age of poultry.

Means egg white weight ranged from 4.93-5.63 g or 54.54 -58.37%. Song et al. (2000) revealed that the standard weight of egg white is approximately 61.2%. The results in this study showed that there was no significant difference ($p>0.05$) with the average egg yolk weight value being R1 2.71 g \pm 0.15, R2 2.65 g \pm 0.32, R3 2.57 g \pm 0.30, R4 2.81 g \pm 0.28. The mean for all treatments was 2.69 g \pm 0.26. Although the results of this study had no effect ($p>0.05$), it was still included in the regular or good category, which was supported by Yuwanta (2010), who stated that the standard weight of quail egg yolks was 2.5-3.4 grams. In Nastiti et al. (2014) research with an average of 3.21 - 3.36 grams. The weight of quail egg yolk did not significantly affect this study due to the type of feed given. The level of consumption of quail feed was the same, and only differences in protein and energy metabolism levels in each treatment (Hanapis, 2020). Tugiyanti and Iriyanti (2012) suggested the influence of yolk weight on eggs, namely, ovarian development, poultry body weight, feed quality and quantity, age at sexual maturity, environment, and disease.

Means percentage value of egg white is R1 55.39 \pm 0.64%, R2 54.54 \pm 1.38%, R3 58.37 \pm 6.05%, R4 57.07 \pm 1.30 %. The mean of all treatments was 56.34 \pm .24%. The average percentage value of egg yolk weight was R1 30.32 \pm 1.20%, R2 29.17 \pm 0.96%, R3 28.61 \pm 1.39%, R4 28.31 \pm 1.60%. The mean of all treatments was 29.10 \pm 1.41%.

Egg white from eggs will decrease because the absorbed nutrients will form egg yolks first and then the formation of egg albumen. Therefore, if the yolk increases, the percentage in white will decrease (Dharmayanti et al., 2019). Furthermore, Suarjana et al. (2018) stated that the percentage of egg whites was negatively correlated with the percentage of yolks, i.e., when the percentage of whites decreased, the percentages of yolks increased. Suprijatna et al. (2005) reported that the egg-making process starts

from the formation and release of the ovum (yolk) and then enters the infundibulum, which will immediately pass through the extended surface of the oviduct from the oviduct.

The results in this study showed that there was no significant difference ($p>0.05$) with the average value of the Haugh unit being R1 82.11 \pm 1.39, R2 84.04 \pm 1.59, R3 84.41 \pm 0.52, R4 84.83 \pm 1.91. The mean of all treatments was 83.85 \pm 1.68. The average of this study is lower than the results of research by Silaban et al. (2019), which used castor bean meal in the ration. The average Haugh unit value produced was 86.72 in control treatment. However, the HU value of the experimental eggs is included in the AAA quality category (HU value is more than 79) (USDA, 2004).

CONCLUSION

This study concluded that a ration with a protein balance of 17% and metabolizable energy of 2800 kcal/kg was able to maintain the internal and external quality of eggs in quail and produce standard egg quality.

CONFLICT OF INTEREST

There was no conflict of interest in this research with other parties, both individuals and organizations.

ACKNOWLEDGMENT

The authors would like to thank the Directorate General of Research and Technology, and Higher Education of Republic of Indonesia for the research grant (No. 09/E1/KPT/2021 Tanggal 01 Februari 2021)

REFERENCES

- Agustini, M.A., G.A.M.K. Dewi, & I.W. Wijana. 2014. Pengaruh imbalanced energi dan protein ransum terhadap kualitas telur ayam kampung umur 20-30 minggu. *Peternakan Tropika* 2(2): 143-152. [Indonesia]
- Ardiansyah, H.R., Sujana, & W. Tanwiriah. 2016. Pengaruh pemberian tingkat protein dalam ransum terhadap kualitas telur puyuh (*Cortunix-cortunix japonica*). *Jurnal Student e-journal* 5(4):1- 10. [Indonesia]

- Dharmayanti, M.R., I.G.N.G. Bidura, & I.A.P. Utami. 2019. Pengaruh ekstrak air daun kunyit (*Curcuma domestica* Val.) melalui air minum terhadap kualitas fisik telur lohmann brown. *Peternakan Tropika* 7(1):253-268. [Indonesia]
- [Ditjen PKH] Ditjen Peternakan Kesehatan Hewan. 2019. Statistik Peternakan dan Kesehatan Hewan. Ditjen Peternakan Kesehatan Hewan. Jakarta. [Indonesia]
- Gheisari, A., H.A. Halaji, G. Maghsoudinegad, M. Toghiani, A. Alibemani, & S.E. Saeid. 2011. Effect of different dietary levels of energy and protein on performance of japanese quails (*Coturnix-coturnix japonica*). *Proceeding of 2nd International Conference on Agricultural and Animal Science*. 156-159.
- Hanapis, E.J. Guntoro, & Aswana. 2020. Pengaruh penggantian sebagian pakan komersil dengan tepung wortel dengan limbah pasar sampai level 12% terhadap berat telur puyuh (*Coturnix-coturnix japonica*). *Stock Peternakan* 2(1):6-25. [Indonesia]
- Kamran, Z., M. Sarwar, M. Nisa, M.A. Nadeem, S. Mahmood, M.E. Babar, & S. Ahmed. 2008. Effect of low protein diets having constant energy-to-protein ratio on performance and carcass characteristics of broiler chickens from one to thirty-five days of age. *Poultry Science* 87(3):468-474.
- Khalil, M.M. 2015. Use of Enzymes to Improve Feed Conversion Efficiency in Japanese Quail Fed a Lapin-based Diet. The University of Western. Australia.
- Lestari, I.D., V.D. Yunianto, & E. Suprjatna. 2015. Pengaruh penggunaan limbah rumput laut (*Gracilaria verrucosa*) terhadap efisiensi penggunaan protein puyuh jantan umur 6-10 minggu. *Animal Agriculture Journal* 4(2): 252-255. [Indonesia]
- Napirah, A., H. Has, L.O. Nafiu, A. Bain, & T. Saili. 2018. Imbangan protein dan energi berbeda dalam ransum puyuh fase grower terhadap konsumsi pakan, pertambahan bobot badan, dan konversi ransum. *Jitro* 2(5):53-57. [Indonesia]
- Nastiti, R.A. Hermana, & R. Mutia. 2014. Penggunaan dedak gandum kasar (wheat bran) sebagai pengganti jagung dengan kombinasi tepung daun mengkudu (*Morinda citrifolia*) untuk menghasilkan telur puyuh sehat rendah kolesterol dan kaya Vitamin A. *Buletin Makanan Ternak* 101(1):1-12. [Indonesia]
- Nugraha, P., Anggraeni, & N. Hanafi. 2018. Pengaruh pemberian tepung jahe dan tepung kunyit pada ransum terhadap kualitas eksternal telur puyuh. *Jurnal Peternakan Nusantara* 4(1):13-18. [Indonesia]
- Panekanan, J.O., J.C. Loing, R. Rorimpandey, & P.O.V. Waleleng. 2013. Analisis keuntungan usaha beternak puyuh di Kecamatan Sonder Kabupaten Minahasa. *J Zooteck* 32(5):1-10. [Indonesia]
- Parizadian, B., Y.J. Ahangari, M.S. Shargh, & A. Sardarzadeh. 2011. Effects of different levels of L-carnitine supplementation on egg quality and blood parameters of laying Japanese quail. *Intl J Poultry Sci* 10 (8):621-625.
- Silaban, E.V., M. Tafsir, & N.D. Hanafi. 2019. Free Choice Feeding on the Quality of Quail Eggs (*Coturnix-coturnix japonica*). *Indonesian Journal of Agricultural Research* 2(2):110-125.
- Song, K.T., S.H. Choi, & H.R. Oh. 2000. A comparison of egg quality of pheasant, chukar, quail and guinea fowl. *Asian-Aus J Anim Sci* 13(7): 986-990.
- Suarjana, I.P., N.W. Siti, & I.G.N.G. Bidura. 2018. Pengaruh pemberian ekstrak air daun mengkudu (*Morinda citrifolia*) melalui air minum terhadap kualitas fisik telur ayam lohmann brown umur 22-30 minggu. *Peternakan Tropika* 6(1):129-139. [Indonesia]
- Sujana, E., A. Asep, S. Iwan, & W. Tuti. 2020. The Egg Characteristics of malon broiler, Japanese quails and their cross. *Biodiversitas* 21(3):889-895.
- Suprijatna, E., U. Atmomarsono, & R. Kartasudjana. 2005. *Ilmu Dasar Ternak Unggas*. Penebar Swadaya. Jakarta. [Indonesia]
- [USDA] United States Department of Agriculture. 2004 *Egg grading Manual*. Agricultural Handbook Number 75. National Supervisor, Shell Eggs Grading Branch, Poultry Programs USDA-AMS STOP 0258 1400 Independence Avenue. Washington, DC.

- Tugiyanti, E. & N. Iriyanti. 2012. Kualitas eksternal telur ayam petelur yang mendapat ransum dengan penambahan tepung ikan fermentasi menggunakan isolat prosedur anti histamin. *Jurnal Aplikasi Teknologi Pangan* 1(2):44-47. [Indonesia]
- Wahju J. 2004. *Ilmu Nutrisi Unggas*. Gajah Mada University Press. Yogyakarta. [Indonesia]
- Yuwanta, T. 2010. *Telur dan Kualitas Telur*. Gajah Mada University Press. Yogyakarta. [Indonesia]

External and Internal Quality of the Quail Eggs Fed Ration with Different Level of Metabolizable Energy and Protein

ORIGINALITY REPORT

15%

SIMILARITY INDEX

PRIMARY SOURCES

| | | |
|---|-------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 1 | ojs.unida.ac.id Internet | 64 words — 2% |
| 2 | olddrji.lbp.world Internet | 27 words — 1% |
| 3 | etd.repository.ugm.ac.id Internet | 24 words — 1% |
| 4 | jisppd.com Internet | 24 words — 1% |
| 5 | www.ajhssr.com Internet | 20 words — 1% |
| 6 | www.slideshare.net Internet | 20 words — 1% |
| 7 | www.jstage.jst.go.jp Internet | 16 words — 1% |
| 8 | Emilia Hanusova, Cyril Hrnčár, Anton Hanus, Marta Oravcová. "Egg traits in Japanese quails", Acta fytotechnica et zootechnica, 2016 Crossref | 14 words — 1% |
| 9 | ejournal2.undip.ac.id | |

Internet

13 words — < 1%

10 mail.scialert.net

Internet

13 words — < 1%

11 biodiversitas.mipa.uns.ac.id

Internet

12 words — < 1%

12 jurnal.pancabudi.ac.id

Internet

11 words — < 1%

13 repository.futminna.edu.ng:8080

Internet

11 words — < 1%

14 A.B. Thasleem, L.S. David. "Effect of starter and finisher feed replacement time on the performance of broiler chickens", *AGRIEAST: Journal of Agricultural Sciences*, 2017

Crossref

10 words — < 1%

15 Noven Hariyani, Siswanto Siswanto, Sri Suharyati, Purnama Edy Santosa. "TOTAL ERITROSIT DAN LEUKOSIT BROILER BETINA SETELAH PEMBERIAN JINTAN HITAM (*Nigella sativa*) SEBAGAI IMUNOMODULATOR DALAM AIR MINUM", *Jurnal Riset dan Inovasi Peternakan (Journal of Research and Innovation of Animals)*, 2020

Crossref

9 words — < 1%

16 Singsen, E. P.. "Nunc Dimittis", *Poultry Science*, 1976.

Crossref

9 words — < 1%

17 academic.oup.com

Internet

9 words — < 1%

18 helvia.uco.es

Internet

| | | |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| | | 9 words — < 1% |
| 19 | oamjms.eu Internet | 9 words — < 1% |
| 20 | Hernandez, F., M. Lopez, S. Martinez, M. D. Megias, P. Catala, and J. Madrid. "Effect of low-protein diets and single sex on production performance, plasma metabolites, digestibility, and nitrogen excretion in 1- to 48-day-old broilers", Poultry Science, 2012. Crossref | 8 words — < 1% |
| 21 | ejournal3.undip.ac.id Internet | 8 words — < 1% |
| 22 | garuda.kemdikbud.go.id Internet | 8 words — < 1% |
| 23 | media.neliti.com Internet | 8 words — < 1% |
| 24 | talenta.usu.ac.id Internet | 8 words — < 1% |
| 25 | www.acarindex.com Internet | 8 words — < 1% |
| 26 | www.neliti.com Internet | 8 words — < 1% |
| 27 | www.scialert.net Internet | 8 words — < 1% |
| 28 | W. Satria, A. E. Harahap, T. Adelina. "Kualitas Telur Puyuh yang Diberikan Ransum dengan | 7 words — < 1% |

Crossref

29 Muarif, Y Wahyudin, D Merdekawati. "Water
quality at silvoaquaculture pond in indramayu
regency", IOP Conference Series: Earth and Environmental
Science, 2019

6 words — < 1%

Crossref

EXCLUDE QUOTES OFF

EXCLUDE SOURCES OFF

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF