

PAPER NAME

Volatilomics for halal and non-halal meat ball authentication using solid-phase microextraction-gas c

AUTHOR

Lia Amalia

WORD COUNT

10149 Words

CHARACTER COUNT

53623 Characters

PAGE COUNT

17 Pages

FILE SIZE

1.4MB

SUBMISSION DATE

Apr 15, 2023 2:17 PM GMT+7

REPORT DATE

Apr 15, 2023 2:18 PM GMT+7

● 18% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 10% Internet database
- 8% Publications database
- Crossref database
- Crossref Posted Content database
- 13% Submitted Works database

● Excluded from Similarity Report


- Bibliographic material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources
- Manually excluded text blocks



King Saud University

Arabian Journal of Chemistry

www.ksu.edu.sa
www.sciencedirect.com



ORIGINAL ARTICLE

Volatilomics for halal and non-halal meatball authentication using solid-phase microextraction–gas chromatography–mass spectrometry



Agy Wirabudi Pranata^{a,b}, Nancy Dewi Yuliana^{a,b,*}, Lia Amalia^{a,b,d},
Noviyan Darmawan^{b,c}

^a Department of Food Science and Technology, IPB University, 16680 Bogor, Indonesia

^b Halal Science Center, IPB University, 16129 Bogor, Indonesia

^c Department of Chemistry, IPB University, 16680 Bogor, Indonesia

^d Food and Nutrition Study Program, Faculty of Halal Food, Djuanda University, 16720 Bogor, Indonesia

Received 27 December 2020; accepted 29 March 2021

Available online 2 April 2021

KEYWORDS

Volatiles;
Metabolomics;
Halal;
Meat;
Adulteration;
PLS-DA

Abstract The adulteration of beef meatballs with wild boar (*Sus scrova*) meat or chicken may be undertaken for economic reasons. This adulteration is a very sensitive issue, particularly for Muslim consumers, as the consumption of wild boar is strictly prohibited by Islamic law. This study aimed to discriminate volatile compounds in meatballs made from beef, chicken, and wild boar and mixtures thereof using solid-phase microextraction–gas chromatography–mass spectrometry (SPME/GC–MS) and multivariate data analysis. SPME is a non-destructive method for the extraction of volatile compounds and does not alter the original chemical composition of the volatiles. A validated partial least squares discriminant analysis (PLS-DA) model with three classes was used to uncover the discriminating volatiles of each type of meatball. The results indicated that β -cymene, 3-methyl-butanol, and 2-pentanol were among the positive discriminating volatiles with the highest variable importance in projection (VIP) values among the chicken meatballs. The highest VIP positive discriminating volatiles in the beef meatballs were 5-ethyl-m-xylene, benzaldehyde, and 3-ethyl-2-methyl-1,3-hexadiene. The mixed meatballs exhibited an interesting profile, with all appearing in the same group as the pure wild boar meatballs. However, the discriminating volatiles derived from a separate PLS-DA model indicated that they contained different compounds. In the pure wild boar meatballs, six compounds (pentanal, 2,6-dimethylcyclohexanone, 1-undecanol, cyclobutanol, 2,4,5-

* Corresponding author at: Department of Food Science and Technology, IPB University, 16680 Bogor, Indonesia.

E-mail address: nancy_dewi@apps.ipb.ac.id (N.D. Yuliana).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

trimethyl-thiazole, and 5-ethyl-3-(3-methyl-5-phenyl pyrazol-1-yl)-1,2,4-triazol-4-amine) were identified as discriminating volatile compounds with the highest VIP values. These compounds were consistently found as significant discriminating volatile compounds in mixture meatballs group although with different VIP value. This research demonstrated that SPME-GC/MS combined with multivariate data analysis was a fast and reliable method for differentiating meatballs made from beef, chicken, and wild boar meat based on their volatile compound contents.

© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Beef meatballs are one of the most popular processed meat products in Indonesia. Since beef prices are rather high, meatballs are often adulterated by mixing beef with cheaper meats, such as wild boar (*Sus scrova*) or chicken, to illegally obtain economic benefits (Guntarti et al., 2017). This adulteration is a disadvantage for consumers, particularly Muslim ones, who are strictly prohibited from consuming wild boar meat. Wild boars are frequently obtained from recreational animal hunting since they are considered pests of plantation crops. They have larger carcass fatness and loin areas, darker meat color, and leaner and less tender meat compared with those of domestic pigs (Sales and Kotrba, 2013). This makes their meat visually more similar to beef.

Fast, sensitive, and affordable analytical methods are necessary to monitor the enforcement of regulations related to meatball consumer protection and support efforts to control the circulation of processed meat products with inappropriate labels. The two most commonly used methods to detect the contamination or adulteration of meat products are polymerase chain reaction (PCR) and enzyme-linked immunosorbent assay (ELISA). PCR is a DNA-based method and has high sensitivity, being able to detect 0.001–0.1 ng DNA from adulterant species (Sultana et al., 2018). Unlike PCR, ELISA uses a specific protein or peptide from each species as a test target. Several other instruments and methods aimed at detecting meat adulterant have also been developed, especially for halal-testing, including liquid chromatography–mass spectrometry (LC-MS) and Fourier transform infrared spectroscopy (FTIR; Kurniawati et al., 2014; Masiri et al., 2016; Rohman et al., 2020; Xu et al., 2012). However, these techniques are not without limitations. In particular, they require rigorous sample preparation and high technical skills, making them less than ideal for routine analysis.

Several previous studies have focused on the profile discrimination of volatiles to assess meat quality. Extracting volatiles from meat can be achieved using several techniques. In one study, 33 volatiles from chicken breast were successfully identified using distillation in dichloromethane (Ayseli et al., 2014). This method has limitations, as it can potentially lead to the loss of thermally unstable compounds. In addition, FTIR coupled with multivariate data analysis has been reported as a rapid and non-destructive yet powerful technique for determining meat types in meatballs (Rahmania et al., 2015). However, the information that can be obtained from FTIR is limited, as the spectra show only absorption bands attributable to the characteristic frequencies of different functional groups. Gas chromatography–mass spectrometry (GC–MS) provides more detailed information, as the spectra

exhibit the specific mass spectrum of the compounds present in the samples, which are eluted at different retention times (Sim et al., 2014).

Solid-phase microextraction–GC–MS (SPME/GC–MS) is a rapid and straightforward technique integrating volatile compound extraction and analysis (Wang et al., 2018). Recently, SPME/GC–MS combined with principal component analysis (PCA) and orthogonal partial least squares discriminant analysis (OPLS-DA) was used to assess compounds related to pork volatiles during storage. This method revealed that ethanol, 2,3-butanediol, and 2-ethyl-1-hexanol can potentially serve as indicators of pork quality during storage (Song et al., 2021). A similar technique was used to study discriminating volatiles of minced beef and pork (Pavlidis et al., 2019) and raw and cooked beef (Wang et al., 2018).

The objective of the present work was to determine discriminating volatiles of beef, chicken, wild boar meat, and meatballs made from mixtures thereof. In Indonesia, consumers commonly buy ready-to-cook meatballs rather than raw ones. These meatballs have to be re-boiled for few minutes before they are consumed. Cases of adulteration occur often with these ready-to-cook meatballs and are difficult to trace because the modified meatballs have similar physical appearances to the unmodified ones. SPME/GC–MS combined with PCA and PLS-DA was here used to analyze discriminating volatiles in different types of meatball samples. We used PCA as a first-pass unsupervised tool in our volatilomics data, whereas sample classification patterns in score-space are the sole basis for further analysis using supervised methods, such as PLS-DA (Worley and Powers, 2016). PCA is commonly used to assess classification patterns within data sets containing unlabeled data. In addition to PCA and PLS-DA, soft independent modeling of class analogy (SIMCA) may also be used. SIMCA is a supervised method used to extract features and obtain classification tasks, according to which the training data are labeled, and the method is then separately applied to each data class. SIMCA has been demonstrated to be a superior method when working with larger data sets, whereas PCA and PLS-DA are more suitable for classification tasks when one has limited access to data (Nejadgholi and Bolic, 2015). PLS-DA is often used in metabolomics research to build predictive classification models and/or discover biomarkers. In PLS-DA, the ideal number of the modeled class is between two and four. When PLS-DA is used to model more than four classes, the classification results may be difficult to interpret (Eriksson et al., 2006).

In this study, PCA was initially used to observe the classification pattern of the meat samples. Once the PCA model had demonstrated satisfactory performance based on cross-validation parameters, a supervised PLS-DA model was devel-

oped. Discriminating volatile selection from the validated PLS-DA was performed based on the variable importance in projection (VIP) value and correlation coefficients.

9. Materials and methods

2.1. Meat sample collection

All samples were collected in December 2019. The wild boar meat samples used in this experiment were obtained from three male wild boars (10–50 kg) originating from Jambi Forest in Sumatera, Indonesia. The meat samples were chopped from the flanks, ribs, and shanks in equal amounts, tightly packed, and then sealed in plastic bags, put in an icebox, and transported to Bogor. The beef samples were obtained from three cattle (Brahman crossbred cattle, 400–550 kg) and taken from the flank only since this is most commonly used by meatball sellers in Indonesia. Three fresh broiler chicken meat samples were obtained from the local slaughterhouse in Bogor. All meat samples were immediately transported to the laboratory and kept in a freezer (−33 °C) prior to analysis.

2.2. Preparation of meatball samples

The meatballs were prepared according to the cooking method commonly used by meatball sellers in Indonesia, though we did not use any spices or taste enhancers, to avoid masking effects. The meat samples were homogenized using a Phillips ProMix hand blender (HR2533), and the meatballs were made by mixing 200 g of minced meat with 20 g of tapioca flour (Cap Pak Tani, Bogor, Indonesia) and 50 g of ice cubes. The dough was then formed by hand into balls with 3–4 cm diameters, put in water at a temperature of 80 °C–100 °C and boiled for 15 mins. The beef, chicken, and wild boar meatballs were each made in triplicate. Similar steps were conducted to prepare the mixed meatballs. The mixed meatballs also contained 200 g of chicken along with wild boar and with beef in the following ratios: 20:80, 40:60, 60:40, and 80:20. In addition, meatballs made of a combination of beef, chicken, and wild boar with a ratio of 40:40:20 were also prepared. Each mixed

meatball was made in duplicate. A total of 31 meatball samples were used in this study (Table 1). For SPME analysis, the meatballs were crushed with a porcelain mortar and pestle. The crushed meatball samples (8 g) were put in a closed headspace vial with 5 mL of distilled water for exposure to SPME fiber.

18. 3. Analysis of volatile compounds

2.3.1. Headspace solid-phase microextraction procedure

The SPME fiber (DVB/CAR/PDMS, Supelco, Bellefonte, PA, USA) was cleaned before use by heating in a GC–MS injector at 250 °C for 5 mins. The pre-extraction process was carried out by placing the clean SPME fibers into the sample headspace in the vial for 10 mins. The vial was put on a heating plate at a constant temperature of 40 °C. The extraction continued for another 30 mins at the same temperature and with the same vial sample and fiber sphere positions as in the pre-extraction process. This procedure was previously described by Pavlidis et al. (2019). The samples were not stirred, and no NaCl was added.

17. 3.2. Gas chromatography–mass spectrometry analysis

GC–MS analysis was carried out by inserting the fiber that had been exposed to the samples (as described in Section 2.3.1) into the GC–MS injection port. Sample injection was carried out in the split mode (split ratio: 1:2) at 250 °C. The separation of the compounds was carried out on a capillary DB-WAX column with 30 × 0.25 mm dimensions and a film thickness of 0.25 µm, (Agilent Technologies, Santa Clara, USA). The oven temperature was maintained at 40 °C for 5 mins and then increased at 4 °C/min until it reached 150 °C. Next, the temperature was further raised to 250 °C (10 °C/min) and held for 5 mins. The interface temperature was set at 280 °C. The mass spectrometer was operated in the electron ionization mode with the electron energy set at 70 eV and a scanning range of 29–350 *m/z* (speed: 4.37 scans/s; gain factor: 1). The ion source and quadrupole analyzer temperatures were set at 230 °C and 150 °C, respectively. This procedure was previously described (Pavlidis et al., 2019) with small differences, includ-

Table 1 Meatball formulation.

| No. | Chicken (g) | Beef (g) | Wild boar (g) | Tapioca (g) | Ice cube (g) | Code | Replication |
|-----|-------------|----------|---------------|-------------|--------------|--------|-------------|
| 1 | 100 | 0 | 0 | 10 | 25 | C | 3 |
| 2 | 0 | 100 | 0 | 10 | 25 | B | 3 |
| 3 | 0 | 0 | 100 | 10 | 25 | W | 3 |
| 4 | 80 | 0 | 20 | 10 | 25 | CW82 | 2 |
| 5 | 60 | 0 | 40 | 10 | 25 | CW64 | 2 |
| 6 | 40 | 0 | 60 | 10 | 25 | CW46 | 2 |
| 7 | 20 | 0 | 80 | 10 | 25 | CW28 | 2 |
| 8 | 0 | 80 | 20 | 10 | 25 | WB82 | 2 |
| 9 | 0 | 60 | 40 | 10 | 25 | WB64 | 2 |
| 10 | 0 | 40 | 60 | 10 | 25 | WB46 | 2 |
| 11 | 0 | 20 | 80 | 10 | 25 | WB28 | 2 |
| 12 | 20 | 40 | 40 | 10 | 25 | WBC244 | 2 |
| 13 | 40 | 20 | 40 | 10 | 25 | WBC424 | 2 |
| 14 | 40 | 40 | 20 | 10 | 25 | WBC442 | 2 |

ing a different column. Here the DB-WAX column was used instead of the HP-5MS column.

23 3.3. Identification of volatile compounds and data pretreatment

The identification of all volatile compound analytes was estimated using the mass spectra along with the built-in NIST MS 14.0 library as a reference. The compounds were then confirmed using the linear retention index (LRI) from the database and previous reports. To determine the LRI of each analyte, 13 homologous series of a n-alkane solution (C10-40, Polyscience, Niles, IL, USA; 5 mg/L) was used in dichloromethane under the same chromatographic conditions as those used for the samples. 13 The LRI was calculated using the following equation, as described elsewhere (Dool and Kratz, 1962):

$$LRI(\text{compound}) = (100 \times n) + (100 \times z) \times \frac{t_r(\text{compound}) - t_r(n)}{t_r(N) - t_r(n)}$$

Here, LRI (compound) is the LRI of the compound, 12 is the retention time, and n and N are the numbers of carbon atoms in the eluting alkanes before and after the product is produced, respectively. Finally, 12 is the discrepancy in the number of carbon atoms in the smaller and larger alkanes.

1 Multivariate data analysis was performed using the SIMCA-P software (v. 16.0, Sartorius-Umetric, Umeå, Sweden). PCA was used to assess the classification patterns among the different types of meat and meatballs. The PCA performance was evaluated based on the value of the predictive coefficient Q^2X . Next, PLS-DA was used to fine-tune the classification pattern obtained from the PCA. PCA and PLS-DA model validations were conducted by cross-validation, response permutation tests. Cross-validation assesses the reproducibility and the predictive power of PCA and PLS-DA models based on R^2 and Q^2 value, respectively. In PLS-DA, R^2Y represents the goodness of fit, whereas Q^2Y is the accuracy of the prediction parameters. In PCA, the same indicator is represented by the R^2X and Q^2 value. Generally, R^2X , Q^2 , R^2Y , and Q^2Y values of at least 0.5 are considered acceptable (Eriksson et al., 2006). In some cases, 0.4 has been considered acceptable (Worley and Powers, 2016). 1 Additionally, response permutation testing was also conducted since sometimes invalid model might have a high cross-validation Q^2 value. In permutation testing, a reliable model should have a significantly larger Q^2 value than Q^2 values generated from random models using the same data set (Worley and Powers, 2012). All of the validation indicators were also calculated 1 using the SIMCA-P software (v. 16.0, Sartorius-Umetric, Umeå, Sweden).

3. Results and discussion

3.1. Volatile profiling by gas chromatography–mass spectrometry

Table 2 presents the volatile compounds with known molecular formulas that were identified in the meatball samples. Only compounds detected in the control samples are listed, and data from the mixed meatballs are not presented. Overall, 150 volatile compounds were found in the chicken meatballs; 148, in the beef meatballs; and 141, in the wild boar meatballs. These volatiles consisted mainly of 26 aldehydes, ketones, alcohols,

acids, esters, aliphatic hydrocarbons, aromatic hydrocarbons, terpenes, and miscellaneous compounds (Table 2 and Fig. 1). They may have been formed as the result of lipid oxidation, the Maillard reaction, interactions between Maillard reaction products and lipid oxidation products, and/or the thermal degradation of thiamine that occurs during the cooking process (Kosowska et al., 2017). The oxidation of lipids produces a wide range of aliphatic compounds, including saturated and unsaturated hydrocarbons, alcohols, aldehydes, ketones, acids, and esters (Ayseli et al., 2014). At the same time, the Maillard reaction may include many heterocyclic compounds including sulfur and nitrogen compounds (Dashdorj et al., 2015). The data in Table 2 indicate that most of the volatiles detected in this study were the result of lipid oxidation (alcohols, hydrocarbons, aldehydes, and ketones), whereas a few were based on thiamin degradation (sulfur compounds) and the Maillard reaction (acetoin and aldehydes). Some compounds might also have resulted from an interaction of these three processes.

The volatile compounds found in each type of meatball were grouped based on their functional groups (Fig. 1). It can be seen that all the meatballs had a similar composition of volatile components. However, compounds from the ketone group were more common in the chicken meatballs, whereas aldehyde compounds were least present in the chicken. In fact, ketones and aldehydes are major contributors to the “chicken-like” scent (Kerler and Grosch, 1997). A number of compounds, including nonane; 3-methyl-, 2,3-octanedione, 4-nonanone, acetophenone, 6-dodecanone, 20 heptanone, 6-methyl-, 2-methyl-3-octanone, 2,5-octanedione 5-hepten-2-one, 6-methyl-, 2-nonanone, 6,7-dodecanedione, 2-undecanone, 1-dodecen-2-one, 3-tridecanone, and nona-3,5-dien-2-one, were detected in the chicken meatballs. A few of the aforementioned compounds, such as 4-nonanone and acetophenone, have previously been identified in boiled chicken (Kerler and Grosch, 1997), and 2,3-octanedione has been found in raw chicken breast (Ayseli et al., 2014).

Only a few previous reports on the composition of the volatile compounds in fresh or boiled wild boar meat were found. Sales and Kotrba (2013) reported that fried wild boar meat contained 48 volatile compounds, including 16 aldehydes, 5 ketones, 6 alcohols, 8 acids, 4 sulfur compounds, 8 pyrazines, 2 furanones, 1 pyrrole, and 3 aromatic compounds. In the present study, 141 volatile compounds were found in the wild boar meatballs, including 2 sulfur compounds, 6 heterocyclics, 20 terpenes, 13 ketones, 3 esters, 3 acids, 29 alcohols, 26 aldehydes, 17 aliphatic hydrocarbons, 17 cyclic aromatic hydrocarbons, and 5 miscellaneous compounds. Alcohol was the most abundant chemical family in the wild boar meatballs. One of the compounds, 1-octen-3-ol, is an important volatile compound and product related to meat fatty acid autoxidation (Mottram, 1998), leading to a mushroom-like scent (Lammers et al., 2009). Other alcohol compounds found in samples of fresh wild boar meat (Sales and Kotrba, 2013) and fried wild boar meat (Lammers et al., 2009) include 1-pentanol, 1-hexanol, 1-heptanol, and 1-octanol.

The volatile compounds detected in the raw and cooked beef were categorized into eight groups: hydrocarbons, alcohols, aldehydes, acids, esters, ketones, furans, and sulfur compounds. The alcohols, acids, and esters were less diverse in the cooked beef as compared with those in the raw beef. By contrast, the aldehydes and ketones were more diverse in the beef after cooking. In particular, the aldehyde diversity increased

Table 2 Volatile compounds identified in beef, chicken, and wild boar meatballs using SPME/GC–MS.

| Volatilomes | LRI | Method Identification ^a | Peak Area ($\times 10^4$) | | |
|--------------------------------------|------|------------------------------------|-----------------------------|----------------|---------------------|
| | | | Chicken Meatballs | Beef Meatballs | Wild Boar Meatballs |
| Acids | | | | | |
| 16 Amino-6-methylbenzoic acid | 1813 | M | 10 | 9.69 | 1.84 |
| 2-Amino-5-methylbenzoic acid | 1843 | M | 3.63 | 8.44 | 292 |
| Caproic acid | 1845 | L | 1.80 | 87 | – |
| Lauric acid | 2066 | M | – | – | 1.98 |
| Alcohols | | | | | |
| 2-Ethylbutanol | 1114 | M | 1.12 | 866 | 862 |
| 2-Butanol, 3-methyl- | 1133 | L | 857 | 360 | 427 |
| 8 Pentanol | 1146 | L | 1.16 | 380 | 137 |
| 2-Ethylcyclobutanol | 1146 | M | – | – | 98 |
| 3-Methylbutanol | 1215 | M | – | 34 | – |
| 2-Pentanol, 4-methyl- | 1275 | L | 1.61 | 311 | 215 |
| 1-Pentanol | 1289 | L | 3.07 | 76 | 1.48 |
| 3-Methyl-3-butenol | 1265 | L | 539 | – | 649 |
| 6-Methyl-2-heptanol | 1302 | L | 4.10 | 221 | 265 |
| 1-Undecanol | 1328 | M | 1.01 | 74 | 1.50 |
| 1-Hexanol | 1350 | L | 402 | 279 | 933 |
| 1-Octanol, 3,7-dimethyl- | 1389 | M | 592 | 494 | – |
| 2-Butoxyethanol | 1404 | L | 281 | 387 | – |
| Ethanol, 2-(dodecyloxy)- | 1440 | M | 93 | 93 | 171 |
| 1-Octanol, 2-butyl- | 1447 | M | 67 | 1.99 | – |
| 1 Octen-3-ol | 1451 | L | 2.35 | – | 2.09 |
| 1-Heptanol | 1461 | L | 1.22 | 550 | 61 |
| 5-Hepten-2-ol, 6-methyl- | 1476 | L | 87 | 762 | 194 |
| 7-Octen-2-ol, 2,6-dimethyl- | 1471 | L | 129 | 30 | 54 |
| 2-Cyclohexen-1-ol | 1511 | L | 98 | – | – |
| 2-Ethylhexanol | 1523 | L | – | 39 | – |
| Cyclobutanol | – | M | 2.23 | 1.74 | 10 |
| Cyclohexanol, 2- <i>tert</i> -butyl- | 1529 | M | 127 | – | – |
| 1-Octanol | 1558 | M | 1.28 | 1.03 | 368 |
| 1-Terpinenol | 1576 | L | 44 | – | 24 |
| Terpinen-4-ol | 1602 | L | 213 | 54 | 135 |
| 1 Cyclooctanol | 1610 | L | 128 | 85 | 52 |
| 2-Octen-1-ol, (E)- | 1613 | L | 141 | 180 | 32 |
| 2-Nonen-1-ol, (E)- | 1620 | M | 230 | 169 | 400 |
| 2-Methyl-1-indanol | 1644 | M | 33 | 6 | – |
| Phenol, 4-(2-propenyl)- | 1694 | M | 176 | – | – |
| 5-Hexen-2-ol | 1784 | M | – | – | 8 |
| Phenol, 3,5-dimethoxy- | 1936 | M | 189 | 273 | 25 |
| 1-Dodecanol | 1982 | L | 211 | 369 | 117 |
| m-Ethylphenol | 1998 | M | 42 | 59 | 139 |
| 2,4-Di- <i>tert</i> -butylphenol | 1994 | M | 601 | 729 | 494 |
| Aldehydes | | | | | |
| Glutaraldehyde | 1073 | M | 3.30 | 2.07 | 259 |
| Pentanal | – | M | 2.48 | 2.22 | 2.98 |
| Hexanal | 1078 | L | 39.8 | 25.1 | 46.3 |
| Heptanal | 1170 | L | 6.78 | 5.21 | 1.02 |
| 5-Methylhexanal | 1182 | M | – | – | 164 |
| Octanal | 1260 | L | 5.71 | 3.52 | 564 |
| 2-Heptenal, (Z)- | 1313 | L | 1.06 | 272 | 2.83 |
| Nonanal | 1369 | L | 9.79 | 1.17 | 1.86 |
| 2-Dodecenal | 1400 | M | 279 | 246 | 612 |
| 2-Undecenal | 1412 | M | 927 | 149 | 62 |
| 2-Octenal, (E)- | 1432 | L | 157 | 100 | 201 |
| Undecanal | 1429 | M | – | 20 | 408 |
| 1 4-Heptadien-1-al | 1480 | L | 466 | 98 | 18 |
| Decanal | 1492 | L | 376 | 492 | 688 |
| Benzaldehyde | 1501 | L | 1.8 | 3.1 | 237 |
| 2-Nonenal, (E)- | 1511 | M | – | 250 | – |
| 2-Tridecenal, (E)- | 1523 | M | – | – | 335 |
| 2-Decenal, (E)- | 1633 | L | 79 | 88 | 96 |

(continued on next page)

Table 2 (continued)

| Volatilomes | LRI | Method Identification ^a | Peak Area ($\times 10^4$) | | |
|-------------------------------------|------|------------------------------------|-----------------------------|----------------|---------------------|
| | | | Chicken Meatballs | Beef Meatballs | Wild Boar Meatballs |
| 1 Octenal, 2-butyl- | 1669 | L | – | 507 | 267 |
| Benzaldehyde, 3-ethyl- | 1688 | M | 1.1 | 1.05 | 425 |
| 2-Dodecenal, (E)- | 1738 | M | 169 | 35 | 50 |
| 2-Undecenal, (E)- | 1746 | L | 27 | 54 | 8 |
| 2,4-Decadienal | 1760 | L | 14 | 8 | 18 |
| 2,4-Decadienal, (E,E)- | 1803 | L | – | – | 27 |
| Butanal, 3-methyl- | – | M | 1.67 | 800 | – |
| Tetradecanal | 1927 | L | 57 | 498 | – |
| Benzaldehyde, 4-pentyl- | 1998 | L | 146 | 158 | 107 |
| Heptadecanal | 2000 | M | 302 | 266 | 47 |
| 17-Octadecenal | 2025 | M | 24 | 124 | 19 |
| Octadecanal | 2015 | M | – | 138 | 253 |
| Aliphatic hydrocarbons | | | | | |
| 5-Ethyl-2-methyloctane | 1053 | L | 2.3 | 1.34 | 613 |
| 1 Undecane, 3-methyl- | 1101 | M | 1.11 | 840 | 2.59 |
| 2,4-Dimethylhexane | 1105 | M | 1.05 | 537 | 406 |
| Undecane, 5-methyl- | 1129 | M | 2.02 | 1.04 | 176 |
| Undecane, 3,4-dimethyl- | 1135 | M | 1.41 | 549 | 133 |
| Dodecane | 1177 | L | 1 | – | 403 |
| 3,5-Dimethylheptane | 1193 | M | 738 | 56 | – |
| 4-Ethylcyclohexene | 1211 | M | – | – | 70 |
| 2,4,6-Trimethyloctane | 1196 | M | 697 | 189 | 265 |
| Tridecane | 1206 | M | – | 58 | – |
| 2-Methyltridecane | 1210 | L | 611 | 141 | – |
| 3,8-Dimethyldecane | 1216 | M | 1.31 | 114 | – |
| 2-Methyldodecane | 1221 | M | 679 | 292 | – |
| 1 2,6-Dimethylundecane | 1234 | M | 461 | 327 | – |
| 3,7-Dimethylnonane | 1252 | M | 971 | 52 | – |
| Undecane, 4,7-dimethyl- | 1273 | M | 1.34 | 280 | 49 |
| 3,6-Dimethyldecane | 1278 | M | 736 | 538 | – |
| Hexane, 2,3,4-trimethyl- | 1286 | M | 1.01 | – | – |
| Hexane, 3-ethyl-4-methyl- | 1284 | M | 886 | 194 | – |
| 1 Undecane, 3,7-dimethyl- | 1348 | M | 168 | – | – |
| 1 Ethyl-2-methyl-1,3-hexadiene | 1387 | M | 532 | 369 | 83 |
| 1-Octene, 3,7-dimethyl- | 1393 | M | – | – | 38 |
| Tetradecane | 1396 | L | 487 | 256 | 53 |
| 1,3-Hexadiene, 3-ethyl | 1396 | M | – | 190 | – |
| 1-Tetradecene | 1425 | L | 150 | 77 | 21 |
| 2-Methyldecane | 1456 | M | 87 | 156 | 1.18 |
| 1-Hexene, 3,5,5-trimethyl- | 1485 | M | 170 | 113 | – |
| 2-Decanone | 1489 | L | 165 | 206 | 64 |
| Pentadecane | 1603 | M | – | 25 | – |
| 1 2-Undecene, 1-ethyl-, (Z)- | 1540 | M | 824 | 1.08 | – |
| 3,5-Dimethyl-1-hexene | 1564 | M | – | – | – |
| Hexadecane | 1596 | L | 88 | 192 | 96 |
| 1-Decene, 3,4-dimethyl- | 1775 | L | – | – | 16 |
| Cyclic aromatic hydrocarbons | | | | | |
| Ethylbenzene | 1111 | L | 4.13 | 2.75 | 1.1 |
| o-Xylene | 1117 | L | 1.95 | 1.02 | – |
| m-Xylene | 1123 | L | 1.76 | 344 | 314 |
| p-Xylene | 1152 | L | 534 | 118 | 80 |
| Furan, 2-pentyl- | 1201 | L | 2.06 | 1.21 | 966 |
| Mesitylene | 1247 | M | 1.26 | 34 | 137 |
| Benzene, 1,2,3-trimethyl- | 1256 | L | 959 | 208 | 588 |
| Cyclooctane, methyl- | 1295 | M | 301 | – | – |
| cis-2-Methylcyclopentanol | 1285 | M | – | 126 | – |
| m-Xylene, 5-ethyl | 1298 | L | 1.44 | 3.12 | 127 |
| Benzene, 2-propenyl- | 1349 | M | 277 | 289 | 83 |
| 2,6-Dimethylcyclohexanone | 1359 | L | 536 | – | 127 |
| Cyclopentane, nonyl- | 1448 | M | – | – | 172 |
| Acridine, 9-methyl- | 1515 | M | 1.30 | 1.05 | 348 |

Table 2 (continued)


| Volatilomes | LRI | Method Identification ^a | Peak Area ($\times 10^4$) | | |
|--|------|------------------------------------|-----------------------------|----------------|---------------------|
| | | | Chicken Meatballs | Beef Meatballs | Wild Boar Meatballs |
| Isopropylcyclohexane | 1550 | M | – | 22 | 301 |
| 4-Ethylbenzaldehyde | 1652 | L | 40 | 20 | 12 |
| Cyclopropane, nonyl | 1658 | M | 61 | 84 | 4 |
| Naphthalene | 1703 | L | 939 | 155 | 550 |
| Azulene | 1731 | M | – | 22 | – |
| Methyl ethyl cyclopentene | 1754 | M | – | 6 | – |
| Butylated Hydroxytoluene | 1910 | L | 57 | 142 | – |
| Cyclodecasiloxane, eicosamethyl- | 2005 | M | – | – | 26 |
| Indole | 2048 | M | 2.79 | 1.74 | 1.18 |
| Esters | | | | | |
| Methyl caprylate | 1354 | L | 416 | 106 | 50 |
| Methyl caprate | 1571 | L | 61 | 90 | – |
| Methyl salicylate | 1749 | L | 29 | – | 6 |
| Methyl palmitate | 1995 | M | 86 | 13 | 303 |
| Heterocyclics | | | | | |
| Thiophene, 2-pentyl- | 1494 | L | 248 | 223 | 51 |
| 1,2,4-Triazol-4-amine phenylpyrazol-1-yl)- | 1546 | M | 380 | 243 | 584 |
| 2-Methyl-2-thiophenecarboxaldehyde | 1765 | L | – | 4 | 61 |
| 2-Methyl-2-thiazole, 2,4,5-trimethyl- | 1867 | L | – | 32 | 698 |
| Tetrahydrothiopyran-4-one | 2006 | M | – | – | 2 |
| Thiophene, 2-butyl-5-ethyl- | 1999 | M | 22 | – | 14 |
| Thiophene, 2-ethyl-5-isopentyl- | 1996 | M | 41 | 11 | – |
| Ketones | | | | | |
| Nonane, 3-methyl- | 1095 | M | 4.17 | 886 | – |
| 6-Dodecanone | 1213 | M | 945 | – | 424 |
| 2-Heptanone, 6-methyl- | 1224 | L | 175 | 168 | 120 |
| 2,3-Octanedione | 1318 | L | 537 | 211 | 747 |
| 2-Methyl-3-octanone | 1323 | L | 202 | – | 240 |
| 2,5-Octanedione | 1330 | L | 1.49 | 138 | 98 |
| 5-Hepten-2-one, 6-methyl- | 1334 | L | 474 | – | 129 |
| 2-Nonanone | 1385 | L | 285 | – | 257 |
| 6,7-Dodecanedione | 1527 | M | 11 | 137 | 195 |
| 4-Nonanone | 1526 | M | 18 | – | – |
| 2-Undecanone | 1576 | L | 11 | – | 11 |
| 11-Dodecen-2-one | 1596 | M | 7 | 27 | 65 |
| Acetophenone | 1630 | L | 313 | 85 | 43 |
| 3-Tridecanone | 1771 | L | 74 | 12 | 19 |
| Nona-3,5-dien-2-one | 1885 | M | 240 | 44 | – |
| γ -Nonalactone | 1999 | L | 85 | 50 | 44 |
| Sulfuric Compounds | | | | | |
| Disulfide, dimethyl | 1068 | L | 1.99 | 1.06 | 439 |
| Disulfide, di- <i>tert</i> -dodecyl | 1316 | M | 729 | 182 | – |
| Dimethyl trisulfide | 1353 | L | 1.85 | 1.79 | 144 |
| Benzothiazole | 1952 | L | 331 | 623 | 70 |
| Terpenes | | | | | |
| o-Cresol | 1142 | M | 1.10 | 604 | 195 |
| 1,4-Cineol | 1159 | L | 1.41 | 570 | 259 |
| D-Limonene | 1165 | L | 1.01 | 744 | 250 |
| γ -Terpinen | 1226 | M | 948 | 123 | – |
| β -Cymene | 1230 | L | 1.95 | 432 | 117 |
| Styrene | 1237 | L | 897 | 67 | 210 |
| o-Cymene | 1240 | L | 1.16 | 228 | 246 |
| 2-Carene | 1251 | M | 683 | 303 | – |
| α -Terpinolene | 1265 | L | 1.08 | 125 | 84 |
| Copaene | 1421 | L | 200 | 111 | 107 |
| p-Cymenene | 1492 | M | – | – | 199 |
| 1,3,8-p-Menthatriene | 1466 | M | 321 | 119 | – |
| (+)-2-Bornanone | 1485 | L | 58 | 120 | 62 |
| L-Camphor | 1688 | M | – | – | 46 |
| Fenchol | 1580 | L | 35 | 154 | 21 |
| β -Terpineol | 1626 | L | – | – | 49 |

(continued on next page)

Table 2 (continued)

| Volatilomes | LRI | Method Identification ^a | Peak Area ($\times 10^4$) | | |
|-------------------------|------|------------------------------------|-----------------------------|----------------|---------------------|
| | | | Chicken Meatballs | Beef Meatballs | Wild Boar Meatballs |
| 3-p-Menthol | 1637 | L | 67 | 104 | 46 |
| dl-Menthol | 1641 | L | – | – | 49 |
| Isoborneol | 1653 | L | 10 | 9 | 15 |
| o-Xylenol | 1998 | M | 56 | 68 | 6 |
| p-Cresol | 2070 | M | 53 | 33 | 114 |
| m-Cresol | 1997 | M | 158 | 71 | 88 |
| cis-Isoeugenol | 1997 | M | 92 | 17 | 202 |
| Miscellaneous | | | | | |
| Acetoin | 1294 | L | 1.21 | 129 | 321 |
| 2-Ethoxyethyl ether | 1418 | M | 136 | 70 | – |
| Benzyl nitrile | 1920 | L | 239 | 127 | – |
| 5-Methyl-2-phenylindole | 2068 | M | – | 28 | 45 |
| 3-Methyl-2-formyl | 2051 | M | 46 | 88 | – |
| p-Vinylguaiacol | 2021 | M | 209 | 203 | 454 |
| Diethyltoluamide | 2090 | M | – | – | 475 |

^a Reliability of identification (L: MS data and RI in agreement with those of authentic compounds; M: MS data in close agreement with the NIST MS library).

**Fig. 1** Composition of the volatile compounds detected in each type of control meatball.

from four compounds before cooking to 20 compounds after cooking, with hexanal being one of the most abundant (Wang et al., 2018). Overall, 30 aldehydes were detected in the beef meatballs, with butanal being the most abundant (Table 2).

3.2. Volatilomics

Volatilomics is a term for volatilome analysis aimed at the detection, characterization, and quantification of volatile metabolites from organics (Lytou et al., 2019). The volatilome is defined as the group of all volatile organic compounds pro-

duced by a living organism (plants, animals, etc.), an ecosystem, or a substrate (such as food), and it includes exogenously derived compounds (organic and inorganic; Lytou et al., 2019). The volatilomic approach has recently been applied in various research fields, for applications such as plant analysis (Lytou et al., 2019) and the discrimination of beef and pork (Pavlidis et al., 2019). Most volatilomics studies have employed GC-MS and electronic nose methods. Multivariate data analysis has also been used to analyze the resulting high-dimensional data. Various multivariate methods can be used to extract information from large amounts of volatilomics data, with PCA, PLS-DA, and OPLS-DA being the most common (Worley and Powers, 2016).

8 To evaluate the meatball classification pattern based on the volatile compound composition, unsupervised PCA was performed on the GC-MS data (chromatographic relative peak area) using unit-variance (UV) scaling. In UV scaling, the scaling weight is $1/sk$, where sk is the standard deviation of parameter k . Thus, in UV scaling, all variables have an equal opportunity to influence the data, making it more objective than other scaling methods (Eriksson et al., 2006). Multiplicative signal correction (MSC) filtering was applied to remove the signal noise. The main objective of using MSC filtering is to remove artifacts and interference that are not correlated to the presence of the target analytes (Eriksson et al., 2006). After excluding three sample outliers (one 100%-chicken meatball sample, one mixed chicken-wild boar meatball sample with a 4:6 ratio, and one mixed chicken-wild boar meatball sample with a 6:4 ratio), a PCA model with three principal components explaining 59.8% of the variation was obtained using Hotelling's T^2 analysis with a 95% confidence interval. PC1, PC2, and PC3 of the PCA explained 43.8%, 9.14%, and 6.81% of the variation, respectively. Only the first two components are presented (Fig. 2). The aforementioned results are consistent with the minimum requirements for the model mentioned by Worley and Powers (2016), which includes $Q^2 = 0.41$. The score plot revealed three distinct groups: beef meatballs (B1-B3), chicken meatballs (C1-C2; C3 was excluded because it was located outside the 95% confidence interval), and meatballs made from 100% wild boar meat or a mixture of wild boar with beef and/or chicken (Fig. 2). Notably, 5 meatballs made from a mixture of wild boar with beef and/or chicken at different ratios were clustered together with the 100%-wild boar meatballs. This may have occurred because of the strong influence of the volatile components present in 10 wild boar meat. The loading plot revealed several volatiles responsible for the three groupings (Fig. 2). For the beef meatball group, the discriminating compounds were 28-amino-5-methylbenzoic acid, 2-amino-6-methylbenzoic acid, benzaldehyde, dimethyl trisulfide, and 1-octanol. For the chicken meatball group, 5-hepten-2-ol, 6-methylnonane, 3-methyl, and 2-ethylbutanol were among those predominant discriminating compounds. Lauric acid, 2,4-di-tert-butyl-phenol, and furan 2-pentyl were the discriminating compounds for the wild boar meatballs and wild boar-containing meatballs.

To obtain a clearer classification pattern, a supervised multivariate data analysis method (PLS-DA) was employed. Further analysis using a supervised method is only recommended when the PCA model for the same set of data has an acceptable predictive coefficient (Q^2 of at least 0.4; Worley and Powers, 2016), which was fulfilled by the aforementioned PCA model. The PLS-DA score plot exhibited better performance than the PCA. The PLS-DA model with three classes (class 1: chicken meatballs, class 2: beef meatballs, and class 3: wild boar and wild boar-beef-chicken meatballs) had good performance, with a cumulative explained variance of $R^2X = 0.69$, $R^2Y = 0.99$, and $Q^2 = 0.84$ (Zhang et al., 2020). Here, the beef and chicken samples were clustered separately from each other. Again, the wild boar meatballs and meatballs made from mixtures of wild boar with chicken and/or beef at different ratios were grouped separately from the beef and chicken samples (Fig. 3).

Further validation with 100 random permutations was performed. As illustrated in Fig. 4, the values of R^2Y (green circles) and Q^2Y (blue squares) from the permuted analysis (bottom-left corner) were lower than the associated initial values (top-right corner), indicating the stability of the model and the absence of overfitting (Song et al., 2021). The p -value for the cross-validated analysis of variance (CV-ANOVA) was less than 0.005 (2.6×10^{-4}), demonstrating the model validity (Eriksson et al., 2008).




Fig. 2 6 CA loading biplot of meatballs made from 100% beef (B1-B3), 100% chicken (C1-C3), 100% wild boar (W1-W3), and mixtures thereof (WB: wild boar-beef, WC: wild boar-chicken, and WBC: wild boar-beef-chicken). The loading biplot illustrates several markers of the beef, chicken, and mixed meatballs. The numbers after the letters represent the percentages of the respective meats. The last number represents the number of replications.




Fig. 3 PLS-DA plot score of meatball samples (C: chicken, B: beef, W: wild boar, WB: wild boar–beef, CW: chicken–wild boar, and WBC: wild boar–beef–chicken). The number represents the ratio of each meat and replication number.

3.3. Potential volatile marker of wild boar meatballs

To elucidate the volatile compounds that serve as markers for each PLS-DA class, a correlation coefficient and the VIP values were used. The compounds that were positively or negatively correlated with the groupings could be determined using the coefficient, whereas the VIP value has only a positive value. Fifteen compounds with both positive and negative correlation values and the largest VIP values were selected from each PLS-DA class (Table 3).

The volatile with the highest VIP value in the chicken meatball class of the PLS-DA was β -cymene (Table 2). We could find no previous report on the occurrence of this compound in fresh or cooked chicken meat. However, a recent review indicated that cymene was found in essential oils, which are often added to poultry feed as natural antibiotics and immune-stimulants (Brenes and Roura, 2010). A previous study also reported that the second-strongest positive compound, 3-methylbutanal (an aldehyde), was detected in thermally processed chicken as a result of the Maillard reaction (Tian et al., 2007). This was also recently reported as one of the volatiles detected in Dezhou braised chicken (Duan et al., 2015) and grilled chicken (Ngamchuachit et al., 2015).

In the beef class, the most robust discriminator was 5-ethylm-xylene. Other discriminating volatiles, including benzaldehyde, octanol, 2-nonenal, and heptanal, have been found among the volatiles isolated from heat-treated beef, and 2-nonenal was also found in processed pork (Dwivedi and Brockmann, 1975). In this study, heptadecanal exhibited a significant contribution as a discriminating volatile in the beef, though it has been previously found in processed pork and ham (Dwivedi and Brockmann, 1975). Dimethyl trisulfide was also one of the potent odorants identified in stewed beef juice (Guth and Grosch, 1994).

In the wild boar and mixture meatballs group, xylene was identified as the strongest discriminator. This compound has previously been detected as a volatile in processed pork and ham (Dwivedi and Brockmann, 1975). The positive volatile with the second-highest VIP value in the wild boar and mixture




Fig. 4 Permutation test for the PLS-DA model for (A) chicken meatballs, (B) beef meatballs, and (C) wild boar and beef–wild boar–chicken meatballs.

Table 3 Fifteen compounds with positive and negative coefficient values with the highest VIP value selected from each PLS-DA class.

| PLS-DA Class 1 (Chicken Meatballs) | | | | | | | |
|------------------------------------|--------------------------------|------|------------------------------|-----|--------------------------------|------|------------------------------|
| No. | Positive compound | VIP | Chemical group | No. | Negative compound | VIP | Chemical group |
| 1 | β -Cymene | 2.08 | terpenes | 1 | Benzaldehyde | 1.95 | aldehydes |
| 2 | Butanal, 3-methyl- | 2.04 | aldehydes | 2 | 3-Ethyl-2-methyl-1,3-hexadiene | 1.74 | aliphatic hydrocarbons |
| 3 | 1-Pentanol | 1.98 | alcohols | 3 | Benzaldehyde, 4-pentyl- | 1.55 | cyclic aromatic hydrocarbons |
| 4 | 2-Pentanol | 1.84 | alcohols | 4 | Undecane, 5,7-dimethyl- | 1.34 | aliphatic hydrocarbons |
| 5 | 3,8-Dimethyldecane | 1.76 | aliphatic hydrocarbon | 5 | Pentanal | 1.27 | aldehydes |
| 6 | Mesitylene | 1.75 | cyclic aromatic hydrocarbons | 6 | 2-Amino-5-methylbenzoic acid | 1.21 | acids |
| 7 | 3-Methyl-3-butenol | 1.74 | alcohols | 7 | 2-Octenal, 2-butyl- | 1.14 | aldehydes |
| 8 | 1,4-Cineol | 1.7 | terpenes | 8 | 5-Hepten-2-ol, 6-methyl- | 1.12 | alcohols |
| 9 | Undecane, 3,4-dimethyl- | 1.69 | aliphatic hydrocarbons | 9 | 1-Dodecanol | 1.08 | alcohols |
| 10 | Tridecane | 1.56 | aliphatic hydrocarbons | 10 | Decanal | 1.04 | aldehydes |
| 11 | 2-Methyltridecane | 1.47 | aliphatic hydrocarbons | 11 | Butylated hydroxytoluene | 0.99 | cyclic aromatic hydrocarbons |
| 12 | 3,5-Dimethylheptane | 1.47 | aliphatic hydrocarbons | 12 | 2-Undecenal | 0.86 | aldehydes |
| 13 | Disulfide, dimethyl | 1.36 | sulfur compounds | 13 | Azulene | 0.85 | cyclic aromatic hydrocarbons |
| 14 | Undecane, 3,4-dimethyl- | 1.3 | aliphatic hydrocarbons | 14 | 2-Decanone | 0.8 | ketones |
| 15 | Styrene | 1.2 | terpenes | 15 | Phenol, 3,5-dimethoxy- | 0.70 | alcohols |
| Class 2 (Beef meatballs) | | | | | | | |
| No. | Positive compound | VIP | Chemical group | No. | Negative compound | VIP | Chemical group |
| 1 | m-Xylene, 5-ethyl | 1.98 | cyclic aromatic compounds | 1 | 1-Pentanol | 1.99 | alcohols |
| 2 | Benzaldehyde | 1.94 | aldehydes | 2 | Mesitylene | 1.75 | cyclic aromatic hydrocarbons |
| 3 | 1-Ethyl-2-methyl-1,3-hexadiene | 1.74 | aliphatic hydrocarbons | 3 | 3-Methyl-3-butenol | 1.74 | alcohols |
| 4 | 1-Octanol, 3,7-dimethyl- | 1.63 | alcohols | 4 | m-Xylene | 1.70 | cyclic aromatic hydrocarbons |
| 5 | Benzaldehyde, 4-pentyl- | 1.54 | aldehydes | 5 | 2-Methyltridecane | 1.47 | aliphatic hydrocarbons |
| 6 | 1-Octanol | 1.42 | alcohols | 6 | p-Xylene | 1.45 | cyclic aromatic hydrocarbons |
| 7 | 1-Octanol, 2-butyl- | 1.40 | alcohols | 7 | Thiazole, 2,4,5-trimethyl- | 1.43 | heterocyclics |
| 8 | 2-Nonenal, (E)- | 1.36 | aldehydes | 8 | Naphthalene | 1.26 | cyclic aromatic hydrocarbons |
| 9 | Dimethyl trisulfide | 1.35 | sulfur compounds | 9 | 5-Hepten-2-one, 6-methyl- | 1.21 | ketones |
| 10 | Undecane, 5,7-dimethyl- | 1.33 | aliphatic hydrocarbons | 10 | Styrene | 1.21 | terpenes |
| 11 | 2-Butoxyethanol | 1.30 | alcohols | 11 | Lauric acid | 1.17 | acids |
| 12 | 5-Hepten-2-ol, 6-methyl- | 1.21 | alcohols | 12 | Terpinen-4-ol | 1.11 | alcohols |
| 13 | 2-Amino-5-methylbenzoic acid | 1.20 | acids | 13 | Methyl palmitate | 1.09 | esters |
| 14 | Heptanal | 1.16 | aldehydes | 14 | 4-Ethyl-o-xylene | 1.03 | cyclic aromatic hydrocarbons |
| 15 | Heptadecanal | 1.16 | aldehydes | 15 | Phenol, 4-(2-propenyl)- | 0.94 | alcohols |

(continued on next page)

Table 3 (continued)

| Class 3 (Wild boar and mixtures) | | | | | | | |
|----------------------------------|---|------|---------------------------|-----|--------------------------|------|------------------------------|
| No. | Positive compound | VIP | Chemical group | No. | Negative compound | VIP | Chemical group |
| 1 | p-Xylene | 1.40 | cyclic aromatic compounds | 1 | β -Cymene | 2.08 | terpenes |
| 2 | Thiazole, 2,4,5-trimethyl- | 1.40 | heterocyclics | 2 | Butanal, 3-methyl- | 2.05 | aldehydes |
| 3 | 6-Methyl-2-heptanol | 1.31 | alcohols | 3 | m-Xylene, 5-ethyl | 1.93 | cyclic aromatic hydrocarbons |
| 4 | Pentanal | 1.27 | aldehydes | 4 | 2-Pentanol | 1.85 | alcohols |
| 5 | 2-Octenal, 2-butyl- | 1.14 | aldehydes | 5 | 3,8-Dimethyldecane | 1.76 | aliphatic hydrocarbons |
| 6 | Terpinen-4-ol | 1.11 | alcohols | 6 | 3-Methyl-3-butenol | 1.74 | alcohols |
| 7 | 4-Ethyl-o-xylene | 1.03 | cyclic aromatic compounds | 7 | o-Xylene | 1.71 | cyclic aromatic hydrocarbons |
| 8 | Indole | 0.95 | cyclic aromatic compounds | 8 | 1,4-Cineol | 1.7 | terpenes |
| 9 | Phenol, 4-(2-propenyl)- | 0.93 | alcohols | 9 | Undecane, 3,4-dimethyl- | 1.69 | aliphatic hydrocarbons |
| 10 | 2-Undecenal | 0.86 | aldehydes | 10 | Undecane, 5-methyl- | 1.63 | aliphatic hydrocarbons |
| 11 | Azulene | 0.80 | cyclic aromatic compounds | 11 | 1-Octanol, 3,7-dimethyl- | 1.63 | alcohols |
| 12 | 5,2,4-Triazol-4-amine, 5-ethyl-3-(3-methyl-5-phenylpyrazol-1-yl)- | 0.80 | heterocyclic compounds | 12 | Tridecane | 1.56 | aliphatic hydrocarbons |
| 13 | Copaene | 0.80 | terpenes | 13 | Disulfide, dimethyl | 1.36 | sulfur compounds |
| 14 | 2-Decanone | 0.79 | ketones | 14 | Dimethyl trisulfide | 1.20 | sulfur compounds |
| 15 | 1-Octen-3-ol | 0.70 | alcohols | 15 | Heptanal | 1.16 | aldehydes |

meatball class was 2,4,5-trimethyl-thiazoles. Thiazoles have been reported as volatiles directly leading to a complex, meaty aroma (Piao et al., 2019). Pentanal has also been identified in cooked Iberian pigs (Estévez et al., 2003) and had a strong positive effect on the discrimination of minced pork from minced beef (Pavlidis et al., 2019).

As previously described, a strong discriminating volatile was assumed to be present in the 100% wild boar meatballs, which was responsible for clustering all the wild boar-containing meatballs in the same group. To address this, a PLS-DA model with three classes was created, including the meatballs made from 100% chicken, 100% beef, and 100% wild boar (Fig. 5). Although a CV-ANOVA indicated that the model was slightly overfitting, the R^2Y and Q^2Y values were 0.99 and 0.88, respectively. A further test with 200 random permutations also indicated an acceptable model (figure not shown). In this PLS-DA, 2-nonanone and pentanal were the two strongest positive discriminating volatiles from the wild boar group. Other discriminating compounds for 100% wild boar meatballs were summarized in Table 4. The compounds were compared with the 15 strongest positive discriminating volatiles when the wild boar meatballs were put in the same group with the meatballs made from a mixture of wild boar with beef and/or chicken at different ratios (Table 3; derived from the first PLS-DA model). As a result, six compounds (pentanal, 2,6-dimethylcyclohexanone, 1-undecanol, cyclobutanol, 2,4,5-trimethyl-thiazole, and 5-ethyl-3-(3-methyl-5-phenyl pyrazol-1-yl)-1,2,4-triazol-4-amine), were found as discriminating volatiles in mixture meatballs but with




Fig. 5 PLS-DA score plot of meatball volatiles' data. Only samples of 100% chicken (class 1; C1, C2, and C3), 100% beef (class 2; B1, B2, and B3), and 100% wild boar (class 3; W1, W2, and W3) are included.

different VIP value (Table 3). However, the strongest pure wild boar meatballs discriminating volatiles (2-nonanone) was not found among volatiles that positively correlate with mixture meatballs grouping. This compound was reported as one of major ketones found in raw pork (Soncin et al., 2007), but there is no reports on its availability in wild boar.

In the further analysis, we excluded 100% wild boar meatballs data to obtain another PLS-DA with 3 classes (100% chicken, 100% beef, and mixture meatballs) Fig. 6. Fifteen

Table 4 List of 15 discriminating volatile compounds in 100% wild boar meatballs with the highest VIP value.

| No. | Positive Compound | VIP | Chemical Group |
|-----|---|------|------------------------------|
| 1 | 2-Nonanone | 1.6 | ketones |
| 2 | Pentanal | 1.47 | aldehydes |
| 3 | 2,6-Dimethylcyclohexanone | 1.34 | cyclic aromatic hydrocarbons |
| 4 | 1-Undecanol | 1.29 | alcohols |
| 5 | Cyclobutanol | 1.28 | alcohols |
| 6 | 1-Hexanol | 1.27 | alcohols |
| 7 | 3-Ethyl-2-methyl-1,3-hexadiene | 1.21 | aliphatic hydrocarbons |
| 8 | Decanal | 1.19 | aldehydes |
| 9 | Thiophene, 2-pentyl- | 1.16 | heterocyclics |
| 10 | 2-Dodecenal | 1.14 | aldehydes |
| 11 | 5-(2,4-Triazol-4-amine, 5-ethyl-3-(3-methyl-5-phenyl pyrazol-1-yl)- | 1.00 | heterocyclic compounds |
| 12 | Lauric acid | 0.98 | acids |
| 13 | 2-Methyldecane | 0.97 | aliphatic hydrocarbons |
| 14 | Thiazole, 2,4,5-trimethyl- | 0.93 | heterocyclics |
| 15 | Copaene | 0.89 | terpenes |

compounds with both positive and negative correlation values and the largest VIP values were selected from each PLS-DA class and summarized in Table 5. Six discriminating volatiles found in pure wild boar meatballs (pentanal, 2,6-dimethylcyclohexanone, 1-undecanol, cyclobutanol, 2,4,5-trimethyl-thiazole, and 5-ethyl-3-(3-methyl-5-phenyl pyrazol-1-yl)-1,2,4-triazol-4-amine) were consistently found as discriminating volatiles of mixture meatballs class in this new PLS-DA model, although with different VIP value. Similarly, the strongest discriminating compound, 2-nonanone, was not found.

These data may partially support the hypothesis that several strong wild boar-discriminating volatiles heavily influenced the clustering of all wild boar-containing meatballs in the same group.

Meat flavor formation during heating, especially when involving volatiles, is a complex process involving various reactions, including the Maillard and unsaturated lipid reactions. Compounds resulting from the Maillard reaction may also react with those from the unsaturated lipid degradation. The exact volatile composition of the meat flavor formed by these reactions depends on not only the types of precursors present in the meat but also the temperature and reaction time (Aaslyng and Meinert, 2017). Meatballs made from a mixture of different types of meat (chicken, beef, and wild boar) at different ratios may develop different volatiles, as concentrations of the precursors vary. This assumption might explain why the discriminatory volatiles of wild boar meatballs were not exactly the same with when clustered together with mixed meatballs as when separated from them.

This study did not include commercial meatballs in its analysis. Instead, the present work is a preliminary study with a very simple meatball formulation. Commercial meatballs typically have a much more complex formulation. Besides meat, flour, salt, and pepper, commercial meatballs may also contain garlic, beef flavor, or a taste enhancer, which could affect the selection of volatile markers. Further research using more complex meatball formulations resembling those of commercial meatballs and including samples of commercial meatballs themselves is required.

In addition, this study did not consider the effect of the animal feed, which can significantly contribute to meat's volatile composition. The volatile compounds in cooked meat can be directly diverted from animal feed into the tissue by the transformation of feed molecules through the action of ruminal microorganisms or by both the Maillard reaction and the oxidation of lipids during the heating process (Vasta and Priolo,




Fig. 6 PLS-DA score plot of meatball volatile compounds data. Only samples of 100% chicken (class 1; C1, C2, and C3), 100% beef (class 2; B1, B2, and B3), and mixture meatballs (class 3; WB with different compositions) are included. Pure wild boar meatballs are excluded. Only the first two PC is presented (PC1 = 46.4%, PC2 16.4%, $R^2Y = 0.984$, $Q^2Y = 0.835$).

Table 5 Fifteen compounds with positive and negative coefficient values with the highest VIP value selected from each class of PLS-DA (100% chicken, 100% beef, and mixture meatballs).

| PLS-DA Class 1 (Chicken meatballs) | | | | | | | |
|------------------------------------|------------------------------|------|------------------------------|-----|---|------|------------------------------|
| No. | Positive Compound | VIP | Chemical Group | No. | Negative Compound | VIP | Chemical Group |
| 1 | Butanal, 3-methyl- | 2.00 | aldehydes | 1 | Benzaldehyde | 1.92 | aldehydes |
| 2 | 1-Pentanol | 1.77 | alcohols | 2 | Ethyl-2-methyl-1,3-hexadiene | 1.49 | aliphatic hydrocarbons |
| 3 | 1,4-Cineol | 1.63 | terpenes | 3 | 2-Undecenal | 1.38 | aldehydes |
| 4 | o-Xylene | 1.60 | cyclic aromatic compounds | 4 | 5-Hepten-2-ol, 6-methyl- | 1.3 | alcohols |
| 5 | 2-Pentanol | 1.54 | alcohols | 5 | Pentanal | 1.2 | aldehydes |
| 6 | Mesitylene | 1.53 | cyclic aromatic hydrocarbons | 6 | 2-Octenal, 2-butyl- | 1.16 | aldehydes |
| 7 | 2-Methyltridecane | 1.52 | aliphatic hydrocarbons | 7 | Undecane, 5,7-dimethyl- | 1.15 | aliphatic hydrocarbons |
| 8 | 3,8-Dimethyldecane | 1.50 | aliphatic hydrocarbon | 8 | Cyclobutanol | 1.12 | alcohols |
| 9 | m-Xylene | 1.49 | cyclic aromatic compounds | 9 | Thiophene, 2-pentyl- | 1.02 | heterocyclics |
| 10 | 3,5-Dimethylheptane | 1.45 | aliphatic hydrocarbons | 10 | 1-Undecanol | 0.89 | alcohols |
| 11 | 3-Methyl-3-butenol | 1.45 | alcohols | 11 | 2,4-Triazol-4-amine, 5-ethyl-3-(3-methyl-5-phenylpyrazol-1-yl)- | 0.88 | heterocyclic compounds |
| 12 | Tridecane | 1.42 | aliphatic hydrocarbons | 12 | Decanal | 0.87 | aldehydes |
| 13 | Undecane, 3,4-dimethyl- | 1.40 | aliphatic hydrocarbons | 13 | Butylated Hydroxytoluene | 0.85 | cyclic aromatic hydrocarbons |
| 14 | Styrene | 1.36 | terpenes | 14 | 2-Decanone | 0.83 | ketones |
| 15 | Disulfide, dimethyl | 1.36 | sulfuric compounds | 15 | 1-Hexanol | 0.79 | alcohols |
| PLS-DA Class 2 (Beef meatballs) | | | | | | | |
| No. | Positive Compound | VIP | Chemical Group | No. | Negative Compound | VIP | Chemical Group |
| 1 | m-Xylene, 5-ethyl | 2.11 | cyclic aromatic compounds | 1 | 1-Pentanol | 1.77 | alcohols |
| 2 | Benzaldehyde | 1.92 | aldehydes | 2 | Mesitylene | 1.53 | cyclic aromatic hydrocarbons |
| 3 | Tetradecanal | 1.84 | aldehydes | 3 | 2-Methyltridecane | 1.52 | aliphatic hydrocarbon |
| 4 | Ethyl-2-methyl-1,3-hexadiene | 1.50 | aliphatic hydrocarbons | 4 | m-Xylene | 1.49 | cyclic aromatic compounds |
| 5 | 1-Octanol, 3,7-dimethyl- | 1.49 | alcohols | 5 | 3,5-Dimethylheptane | 1.49 | aliphatic hydrocarbons |
| 6 | 1-Octanol, 2-butyl- | 1.48 | alcohols | 6 | 3-Methyl-3-butenol | 1.45 | alcohols |
| 7 | 2-Amino-5-methylbenzoic acid | 1.42 | acids | 7 | Styrene | 1.36 | terpenes |
| 8 | 2-Ethylbutanol | 1.35 | alcohols | 8 | p-Xylene | 1.35 | cyclic aromatic |
| 9 | 5-Hepten-2-ol, 6-methyl- | 1.3 | alcohols | 9 | Nonanal | 1.29 | aldehydes |
| 10 | 1-Octanol | 1.24 | alcohols | 10 | Thiazole, 2,4,5-trimethyl- | 1.28 | heterocyclics |
| 11 | Heptanal | 1.19 | aldehydes | 11 | 6-Dodecanone | 1.27 | ketones |
| 12 | Benzaldehyde, 4-pentyl- | 1.17 | aldehydes | 12 | 5-Hepten-2-one, 6-methyl- | 1.19 | ketones |
| 13 | Undecane, 5,7-dimethyl- | 1.15 | aliphatic hydrocarbons | 13 | Lauric acid | 1.08 | acids |
| 14 | Dimethyl trisulfide | 1.09 | sulfuric compounds | 14 | Terpinen-4-ol | 1.02 | alcohols |
| 15 | 2-Nonenal, (E)- | 1.08 | aldehydes | 15 | Methyl palmitate | 1.01 | esters |
| PLS-DA Class 3 (Mixtures) | | | | | | | |
| No. | Positive Compound | VIP | Chemical Group | No. | Negative Compound | VIP | Chemical Group |
| 1 | 2-Undecenal | 1.38 | aldehydes | 1 | m-Xylene, 5-ethyl | 2.11 | cyclic aromatic compounds |
| 2 | p-Xylene | 1.35 | cyclic aromatic compounds | 2 | Butanal, 3-methyl- | 1.99 | aldehydes |
| 3 | Thiazole, 2,4,5-trimethyl- | 1.26 | heterocyclics | 3 | Benzaldehyde | 1.92 | aldehydes |
| 4 | Pentanal | 1.2 | aldehydes | 4 | 1,4-Cineol | 1.63 | terpenes |

Table 5 (continued)

| PLS-DA Class 1 (Chicken meatballs) | | | | | | | |
|------------------------------------|---|------|------------------------------|-----|--------------------------|------|------------------------------|
| No. | Positive Compound | VIP | Chemical Group | No. | Negative Compound | VIP | Chemical Group |
| 5 | 2-Octenal, 2-butyl- | 1.16 | aldehydes | 5 | o-Xylene | 1.6 | cyclic aromatic compounds |
| 6 | Cyclobutanol | 1.12 | alcohols | 6 | 2-Pentanol | 1.54 | alcohols |
| 7 | 2,6-Dimethylcyclohexanone | 1.05 | cyclic aromatic hydrocarbons | 7 | 1-Octanol, 3,7-dimethyl- | 1.49 | alcohols |
| 8 | 2,4-Heptadien-1-al | 1.04 | aldehydes | 8 | 5-Ethyl-2-methyloctane | 1.48 | aliphatic hydrocarbons |
| 9 | Cyclohexanol, 2- <i>tert</i> -butyl- | 1.03 | alcohols | 9 | 3-Methyl-3-butenol | 1.46 | alcohols |
| 10 | 1-Octen-3-ol | 0.97 | alcohols | 10 | Undecane, 5-methyl- | 1.45 | aliphatic hydrocarbons |
| 11 | Phenol, 4-(2-propenyl)- | 0.96 | alcohols | 11 | Undecane, 3,4-dimethyl- | 1.40 | aliphatic hydrocarbons |
| 12 | 1-Undecanol | 0.88 | alcohols | 12 | Ethylbenzene | 1.40 | cyclic aromatic hydrocarbons |
| 13 | 2,4-Triazol-4-amine, 5-ethyl-3-(3-methyl-5-phenylpyrazol-1-yl)- | 0.88 | heterocyclics | 13 | Disulfide, dimethyl | 1.36 | sulfuric compounds |
| 14 | Caproic acid | 0.86 | acids | 14 | 2-Ethylbutanol | 1.35 | alcohols |
| 15 | Indole | 0.86 | cyclic aromatic hydrocarbons | 15 | 2-Butoxyethanol | 1.20 | alcohols |

2006). An example of such a case is in a study by (Resconi et al., 2010), in which male Corriedale lambs that were only fed by pasture were found to have significantly lower levels of alkanals, alkadienals, and ketones compared with those of lambs fed by pasture and concentrate and concentrate-plus-lucerne hay.

4. Conclusion

This study revealed that it is possible to classify meatball products according to the different types of meat they contain based on volatile profiles, including halal (beef and chicken) and non-halal species (wild boar). The PLS-DA model with three classes indicated that β -cymene, 3-methyl-butanol, and 2-pentanol were among the positive discriminating volatiles with the highest VIP in the chicken meatball group, whereas benzaldehyde, 2-ethyl-2-methyl-1,3-hexadiene, and 4-pentyl-benzaldehyde were the three strongest negative discriminating volatiles in this group. In the beef meatball class, the highest VIP positive discriminating volatiles were 5-ethyl-m-xylene, benzaldehyde, and 3-ethyl-2-methyl-1,3-hexadiene, whereas the three highest VIP negative ones were 1-pentanol, mesitylene, and 3-methyl-3-butenol. The mixed meatballs exhibited an interesting profile with all being clustered with the 100%-wild boar meatballs. Discriminating volatiles derived from a separate PLS-DA model pointed to a consistent 6 compounds, those are 2-pentanol, 2,6-dimethylcyclohexanone, 1-undecanol, cyclobutanol, 2,4,5-trimethyl-thiazole, and 5-ethyl-3-(3-methyl-5-phenyl pyrazol-1-yl)-1,2,4-triazol-4-amine. These compounds were identified as significant discriminating compounds in pure wild boar meatballs and mixture meatballs, but with different VIP value in each PLS-DA models. Further study to link the volatile characteristics of each class with the respective aroma perceptions using gas chromatography-olfactometry (GC-O) is recommended.

9 Funding

This study was supported by the Ministry of Research and Technology/National Agency for Research and Innovation, Republic of Indonesia, through the Penelitian Dasar Unggulan Perguruan Tinggi 2020 scheme under contract number 1/AMD/E1/KP.PTNBH/2020.

CRedit authorship contribution statement

Agy Wirabumi Pranata¹⁰ investigation, Formal analysis, Writing - original draft. Nancy Dewi Yuliana: Conceptualization, Methodology, Funding acquisition, Writing - review & editing, Supervision. Lia Amalia: Formal analysis, Project administration. Noviyan Darmawan: Validation, Visualization, Writing - review & editing, Supervision.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Aaslyng, M.D., Meinert, L., 2017. Meat flavour in pork and beef – from animal to meal. *Meat Sci.* 132, 112–117. <https://doi.org/10.1016/j.meatsci.2017.04.012>.
- Ayseli, M.T., Filik, G., Selli, S., 2014. Evaluation of volatile compounds in chicken breast meat using simultaneous distillation and extraction with odour activity value. *J. Food Nutr. Res.* 53, 137–142.
- Brenes, A., Roura, E., 2010. Essential oils in poultry nutrition: main effects and modes of action. *Anim. Feed Sci. Technol.* 158 (1-2), 1–14. <https://doi.org/10.1016/j.anifeedsci.2010.03.007>.

- Dashdorj, D., Amna, T., Hwang, I., 2015. Influence of specific taste-active components on meat flavor as affected by intrinsic and extrinsic factors: an overview. *Eur. Food Res. Technol.* 241 (2), 157–171. <https://doi.org/10.1007/s00217-015-2449-3>.
- van den Dool, H., Kratz, P.D., 1962. A generalization of the retention index system including linear temperature programmes gas-liquid partition chromatography. *J. Chromatogr.* 11, 461–471. https://doi.org/10.1007/978-3-319-70262-9_7.
- Duan, Y., Zheng, F., Chen, H., Huang, M., Xie, J., Chen, F., Sun, B., 2015. Analysis of volatiles in Dezhou Braised Chicken by comprehensive two-dimensional gas chromatography/high resolution-time of flight mass spectrometry. *LWT – Food Sci. Technol.* 60 (2), 1235–1242. <https://doi.org/10.1016/j.lwt.2014.09.006>.
- Dwivedi, B.K., Brockmann, M.C., 1975. Meat flavor. *CRC Crit. Rev. Food Technol.* 5, 487–535. <https://doi.org/10.1080/10408397509527183>.
- Eriksson, L., Johansson, E., Kettenah-Wold, N., Trygg, J., Wikström, C., Wold, S., 2006. Multi- and Megavariate Data Analysis. Umetrics AB, Umeå.
- Eriksson, L., Trygg, J., Wold, S., 2008. CV-ANOVA for significance testing of PLS and OPLS models. *J. Chemom.* 22, 594–600. <https://doi.org/10.1002/cem.1187>.
- Estévez, M., Morcuende, D., Ventanas, S., Cava, R., 2003. Analysis of volatiles in meat from Iberian pigs and lean pigs after refrigeration and cooking by using SPME-GC-MS. *J. Agric. Food Chem.* 51 (11), 3429–3435. <https://doi.org/10.1021/jf026218h>.
- Guntarti, A., Martono, S., Yuswanto, A., Rohman, A., 2017. Analysis of beef meatball adulteration with wild boar meat using real-time polymerase chain reaction. *Int. Food Res. J.* 24, 2451–2455.
- Guth, H., Grosch, W., 1994. Identification of the character impact odorants of stewed beef juice by instrumental analyses and sensory studies. *J. Agric. Food Chem.* 42 (12), 2862–2866. <https://doi.org/10.1021/jf00048a039>.
- Kerler, J., Grosch, W., 1997. Character impact odorants of boiled chicken: changes during refrigerated storage and reheating. *Eur. Food Res. Technol.* 205 (3), 232–238. <https://doi.org/10.1007/s002170050157>.
- Kosowska, M., Majcher, M.A., Fortuna, T., 2017. Volatile compounds in meat and meat products. *Food Sci. Technol.* 37, 1–7. <https://doi.org/10.1590/1678-457X.08416>.
- Kurniawati, E., Rohman, A., Triyana, K., 2014. Analysis of lard in meatball broth using Fourier transform infrared spectroscopy and chemometrics. *Meat Sci.* 96 (1), 94–98. <https://doi.org/10.1016/j.meatsci.2013.07.003>.
- Lammers, M., Dietze, K., Ternes, W., 2009. A comparison of the volatile profiles of frying European and Australian wild boar meat with industrial genotype pork by dynamic headspace-GC/MS analysis. *J. Muscle Foods* 20, 255–274. <https://doi.org/10.1111/j.1745-4573.2009.00146.x>.
- Lytou, A.E., Panagou, E.Z., Nychas, G.J.E., 2019. Volatilomics for food quality and authentication. *Curr. Opin. Food Sci.* 28, 88–95. <https://doi.org/10.1016/j.cofs.2019.10.003>.
- Masiri, J., Benoit, L., Barrios-Lopez, B., Thienes, C., Meshgi, M., Agapov, A., Dobritsa, A., Nadala, C., Samadpour, M., 2016. Development and validation of a rapid test system for detection of pork meat and collagen residues. *Meat Sci.* 121, 397–402. <https://doi.org/10.1016/j.meatsci.2016.07.006>.
- Mottram, D.S., 1998. Flavour formation in meat and meat products: a review. *Food Chem.* 62 (4), 415–424. [https://doi.org/10.1016/S0308-8146\(98\)00076-4](https://doi.org/10.1016/S0308-8146(98)00076-4).
- Nejadgholi, I., Bolic, M., 2015. A comparative study of PCA, SIMCA and Cole model for classification of bioimpedance spectroscopy measurements. *Comput. Biol. Med.* 63, 42–51. <https://doi.org/10.1016/j.combiomed.2015.05.004>.
- Ngamchuachit, P., Kitai, Y., Keeratipibul, S., Phuwapraisirisan, P., 2015. View of comparison of dynamic headspace trapping on Tenax TA and eadspace stir bar sorptive extraction for analysis of vrilled Chicken (Yakitori) Volatiles. *LWT – Food Sci. Technol.* 60, 1235–1242.
- Pavlidis, D.E., Mallouchos, A., Ercolini, D., Panagou, E.Z., Nychas, G.J.E., 2019. A volatilomics approach for off-line discrimination of minced beef and pork meat and their admixture using HS-SPME GC/MS in tandem with multivariate data analysis. *Meat Sci.* 151, 43–53. <https://doi.org/10.1016/j.meatsci.2019.01.003>.
- M.Y. Piao H.J. Lee H.I. Yong S.-H. Beak H.J. Kim C. Jo K.G. Wiryawan M. Baik Comparison of reducing sugar content, sensory traits, and fatty acids and volatile compound profiles of the longissimus thoracis among Korean cattle, Holsteins, and Angus steers Asian-Australasian J. Anim. Sci. 32 1 2019 126 136 10.5713/ajas.18.0065
- Rahmania, H., Sudjadi, A., Rohman, A., 2015. The employment of FTIR spectroscopy in combination with chemometrics for analysis of rat meat in meatball formulation. *Meat Sci.* 100, 301–305. <https://doi.org/10.1016/J.MEATSCI.2014.10.028>.
- Resconi, Virginia C, Campo, M Mar, Montossi, Fabio, Ferreira, Vicente, Sañudo, Carlos, Escudero, Ana, 2010. Relationship between odour-active compounds and flavour perception in meat from lambs fed different diets. *Meat Science* 85, 700–706. <https://doi.org/10.1016/j.meatsci.2010.03.027>.
- Rohman, A., Windarsih, A., Erwanto, Y., Zakaria, Z., 2020. Review on analytical methods for analysis of porcine gelatine in food and pharmaceutical products for halal authentication. *Trends Food Sci. Technol.* 101, 122–132. <https://doi.org/10.1016/j.tifs.2020.05.008>.
- Sales, James, Kotrba, Radim, 2013. Meat from wild boar (*Sus scrofa* L.): a review. *Meat Sci.* 94 (2), 187–201. <https://doi.org/10.1016/j.meatsci.2013.01.012>.
- Sim, Siong Fong, Lee, Terri Zhuan Ean, Mohd Irwan Lu, Nurul Aida Lu, Samling, Benedict, 2014. Synchronized analysis of FTIR spectra and GCMS chromatograms for evaluation of the thermally degraded vegetable oils. *J. Anal. Methods Chem.* 2014, 1–9. <https://doi.org/10.1155/2014/271970>.
- Soncin, S., Chiesa, L.M., Cantoni, C., Biondi, P.A., 2007. Preliminary study of the volatile fraction in the raw meat of pork, duck and goose. *J. Food Compos. Anal.* 20 (5), 436–439. <https://doi.org/10.1016/j.jfca.2006.09.001>.
- Song, Xuechao, Canellas, Elena, Nerin, Cristina, 2021. Screening of volatile decay markers of minced pork by headspace-solid phase microextraction-gas chromatography-mass spectrometry and chemometrics. *Food Chem.* 342, 128341. <https://doi.org/10.1016/j.foodchem.2020.128341>.
- Sultana, Sharmin, Hossain, MA Motalib, Zaidul, ISM, Ali, M Eaqub, 2018. Multiplex PCR to discriminate bovine, porcine, and fish DNA in gelatin and confectionery products. *LWT.* 92, 169–176. <https://doi.org/10.1016/j.lwt.2018.02.019>.
- Tian, Hong Yu, Zhang, Jie, Sun, Bao Guo, Huang, Ming Quan, Li, Jian Rong, Han, Xiao Xiang, 2007. Preparation of natural isovaleraldehyde by the Maillard reaction. *Chinese Chem. Lett.* 18 (9), 1049–1052. <https://doi.org/10.1016/j.ccllet.2007.07.009>.
- Vasta, V., Priolo, A., 2006. Ruminant fat volatiles as affected by diet. A review. *Meat Sci.* 73 (2), 218–228. <https://doi.org/10.1016/j.meatsci.2005.11.017>.
- Wang, Xiaodan, Zhu, Lingtao, Han, Yunxiu, Xu, Liping, Jin, Jiahui, Cai, Yingming, Wang, Hongmei, 2018. Analysis of volatile compounds between raw and cooked beef by HS-SPME-GC-MS. *J. Food Process. Preserv.* 42 (2), e13503. <https://doi.org/10.1111/jfpp.13503>.
- Worley, Bradley, Powers, Robert, 2016. PCA as a practical indicator of OPLS-DA model reliability. *Curr. Metabolomics* 4 (2), 97–103.
- Worley, B., Powers, R., 2012. Multivariate analysis in metabolomics. *Curr. Metabolomics* 1, 92–107. <https://doi.org/10.2174/2213235X11301010092>.

Xu, L., Cai, C.B., Cui, H.F., Ye, Z.H., Yu, X.P., 2012. Rapid discrimination of pork in Halal and non-Halal Chinese ham sausages by Fourier transform infrared (FTIR) spectroscopy and chemometrics. *Meat Sci.* 92 (4), 506–510. <https://doi.org/10.1016/j.meatsci.2012.05.019>.

Zhang, Jialin, Chen, Hui, Fan, Chunlin, Gao, Shuai, Zhang, Zijuan, Bo, Lin, 2020. Classification of the botanical and geographical origins of Chinese honey based on ¹H NMR profile with chemometrics. *Food Res. Int.* 137, 109714. <https://doi.org/10.1016/j.foodres.2020.109714>.

● **18% Overall Similarity**

Top sources found in the following databases:

- 10% Internet database
- Crossref database
- 13% Submitted Works database
- 8% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

| | | |
|----------|--|---------------|
| 1 | Sriwijaya University on 2022-02-03 Submitted works | 6% |
| 2 | researchgate.net Internet | 2% |
| 3 | rodin.uca.es Internet | 1% |
| 4 | ukm.my Internet | 1% |
| 5 | Lia Amalia, Nancy Dewi Yuliana, Purwantiningsih Sugita, Desi Arofah et... Crossref | <1% |
| 6 | repository.ipb.ac.id:8080 Internet | <1% |
| 7 | dspace.aua.gr Internet | <1% |
| 8 | Sriwijaya University on 2021-10-26 Submitted works | <1% |

| | | |
|----|--|-----|
| 9 | mdpi.com Internet | <1% |
| 10 | Sriwijaya University on 2021-11-03 Submitted works | <1% |
| 11 | dce.uni-sofia.bg Internet | <1% |
| 12 | researchrepository.murdoch.edu.au Internet | <1% |
| 13 | Fitra Tunnisa, Didah Nur Faridah, Ani Afriyanti, Dian Rosalina, Mohama... Crossref | <1% |
| 14 | mdpi-res.com Internet | <1% |
| 15 | repository.ipb.ac.id Internet | <1% |
| 16 | Sriwijaya University on 2022-08-09 Submitted works | <1% |
| 17 | doc-developpement-durable.org Internet | <1% |
| 18 | res.mdpi.com Internet | <1% |
| 19 | Sriwijaya University on 2022-02-03 Submitted works | <1% |
| 20 | University of Technology, Sydney on 2013-11-10 Submitted works | <1% |

| | | | |
|----|--|-----------------|-----|
| 21 | labs.mcdb.ucsb.edu | Internet | <1% |
| 22 | Fereidoon Shahidi, Leon J. Rubin, Lorraine A. D'Souza, Roy Teranishi, R... | Crossref | <1% |
| 23 | Dimitrios E. Pavlidis, Athanasios Mallouchos, Danilo Ercolini, Efstathio... | Crossref | <1% |
| 24 | Rutgers University, New Brunswick on 2019-06-02 | Submitted works | <1% |
| 25 | Sriwijaya University on 2022-02-23 | Submitted works | <1% |
| 26 | University of Lincoln on 2021-10-08 | Submitted works | <1% |
| 27 | downloads.hindawi.com | Internet | <1% |
| 28 | patentimages.storage.googleapis.com | Internet | <1% |

● Excluded from Similarity Report

- Bibliographic material
- Small Matches (Less than 10 words)
- Manually excluded text blocks
- Cited material
- Manually excluded sources

EXCLUDED SOURCES

Agy Wirabudi Pranata, Nancy Dewi Yuliana, Lia Amalia, Noviyana Darmawan. "V... 48%
Crossref

EXCLUDED TEXT BLOCKS

Volatilomics for halal and non-halal meatball authentication using solid-phase micr...
www.researchgate.net

The adulteration of beef meatballs with wild boar (*Sus scrova*) meat or chicken ma...
www.researchgate.net

trimethyl-thiazole, and 5-ethyl-3-(3-methyl-5-phenyl pyrazol-1-yl)-1,2,4-triazol-4-a...
www.researchgate.net